



TEMPERATURE AND AQUATIC LIFE

LABORATORY INVESTIGATIONS SERIES

TECHNICAL ADVISORY AND INVESTIGATIONS BRANCH
FEDERAL WATER POLLUTION CONTROL ADMINISTRATION
UNITED STATES DEPARTMENT OF THE INTERIOR
5555 Ridge Avenue
Cincinnati, Ohio 45213



FOREWORD

The Laboratory Investigations series was initiated by the Technical Advisory and Investigations Branch in 1963. The series was planned to describe laboratory methods and techniques and to disseminate information that may be of interest and use to other activities of FWPCA.

The current addition to the series is a literature review of the effects of thermal pollution on the aquatic ecosystem. Thermal pollution is a rapidly increasing problem and this review will aid in evaluating existing problems and in the prevention of future problems.

December 1967

TEMPERATURE AND AQUATIC LIFE

Laboratory Investigations - Number 6

Technical Advisory and Investigations Branch
Technical Services Program
Federal Water Pollution Control Administration
United States Department of the Interior
Cincinnati, Ohio

December 1967

CHEMICAL REACTIONS JAMES L. HOLDAWAY
Chemist

BACTERIA LOUIS A. RESI
Microbiologist

FRESHWATER FISHES NELSON A. THOMAS
Aquatic Biologist

MARINE, ESTUARINE AND ANADROMOUS FISHES . . . LOYS P. PARRISH
Aquatic Biologist

AQUATIC PLANTS AND BENTHOS R. KEITH STEWART
Aquatic Biologist

EDITOR KENNETH M. MACKENTHUN
Supervisory Aquatic
Biologist

SUBJECT INDEX

	<u>Page</u>
CHEMICAL REACTIONS	1
Ammonia	12
Biochemical oxygen demand	8
Carbon dioxide	10
Dissolved Minerals	14
Henry's law	1
Hydrogen	8
Hydrogen sulfide	9
Methane	7
Nitrogen	7
Oxygen	5
Sulfur dioxide	13
Summary	16
References	17
BACTERIA	19
Bacterial indicators	26
Bacterial survival	25
Microbiotic cycles	21
Self-purification	23
Slime growths	26
Summary	27
References	29

	Page
FRESHWATER FISHES	30
Acclimation.	32
Benefits of thermal pollution.	46
Effects on toxicities.	45
General temperature effects.	30
Maximum temperatures	34
Preferred temperatures	38
Sudden temperature changes	31
Summary.	47
References	49
MARINE ESTUARINE AND ANADROMOUS FISHES.	52
Acclimation.	56
Anadromous fishes	
Eggs.	67
Young	70
Adults.	73
Development.	54
Distribution	55
Ecology.	55
General temperature effects.	52
Marine and estuarine fishes	
Eggs.	57
Young	58
Adults.	63
Metabolism	53
Physiology	53
Reproduction	54

	Page
Tolerance.	57
Summary	75
References	77
AQUATIC PLANTS AND BENTHOS.	83
Bottom organisms	87
Fresh-water algae and other aquatic plants	85
General.	83
Summary.	93
References	94
SELECTED BIBLIOGRAPHY	97
INTRODUCTION.	iv
SUMMARY	v

INTRODUCTION

Temperature, a catalyst, a depressant, an activator, a restrictor, a stimulator, a controller, a killer, is one of the most important and most influential water quality characteristics to life in water. Temperature determines those species that may be present; it activates the hatching of young, regulates their activity and stimulates or suppresses their growth and development; it attracts, and kills when the water becomes too hot or becomes chilled too suddenly. Colder water generally suppresses development; warmer water generally accelerates activity and may be a primary cause of aquatic plant nuisances when other environmental factors are suitable.

Because of the importance of this single environmental facet to aquatic ecology, this report was developed to consider some of the features of temperature and its interrelationships. It is divided into five segments; these are:

- I. Chemical Reactions
- II. Bacteria
- III. Freshwater Fishes
- IV. Marine, Estuarine and Anadromous Fishes
- V. Aquatic Plants and Benthos

An extensive temperature bibliography is appended.

SUMMARY

1. Chemical reaction rates vary with temperature, generally increasing as the temperature is increased.
2. The solubility of gases in water varies with temperature. Dissolved oxygen is decreased by the decay or decomposition of dissolved organic substances; the decay rate increases as the temperature of the water increases reaching a maximum at about 30°C (86°F).
3. The temperature of stream water, even during the summer, is below the optimum for pollution-associated bacteria. Increasing the water temperature increases the bacterial multiplication rate when the environment is favorable, and the food supply is abundant. Increasing the water temperature within the growth range of the bacteria causes a more rapid die-off when the food supply is limiting.
4. Warm water fish can survive temporarily in waters heated artificially to 33.9°C (93°F); some fish populations, such as roach, perch, and carp, are reduced at these high temperatures. In cold weather, stream temperatures should be substantially below 33.9°C (93°F) to prevent mortalities when fish move through excessive temperature gradients. Cold water non-anadromous fish populations such as trout should not be subjected to temperatures exceeding 14.5°C (58°F). In cold weather stream temperatures should be below 14.5°C (58°F) to prevent mortalities of cold water fishes.

5. Sudden changes in temperature can be more harmful to some species of fish than continued exposure to a higher temperature.
6. Fish can adapt to higher temperatures faster than to lower temperatures.
7. The maximum temperature for a given species of fish varies with the fish's rate of heating, size, and physiological condition.
8. Fish may starve at elevated temperatures because of their inability to capture food.
9. Fish seek out a preferred temperature at which they can best survive, which is several degrees below their lethal temperature.
10. The toxic effects to fish of certain material increase with temperature.
11. Temperature changes are most important to fish in enclosed areas in the marine environment such as estuaries and bays as opposed to open areas although tolerance to temperature fluctuations is greater in fresh-water and estuarine forms than in open water marine species.
12. There are restricted ranges of temperature within which fish can reproduce successfully; larval development especially requires narrow ranges of temperature. A fish population may exist in a heated area only by continued immigration from the outside. Fish may be absent from such areas during warm summer months and present in cold winter months.
13. Increased temperatures may block the migrations of anadromous fish.
14. Cold is as important to fish populations as heat because of the inability of fish to acclimate quickly to rapid decreases in temperature. Thus, in some areas fish populations may be limited by decreases as well as

increases in temperature. The growth rate of fishes is reduced in waters colder than the optimum temperature range for the species.

15. When water temperatures increase, the predominate algal species change from diatom to green algae and finally at high temperatures to blue-green algae.
16. The number and distribution of bottom organisms decrease as water temperatures increase above 90°F, which is close to the tolerance limit for a "balanced" population. The adult stage of many species is able to tolerate higher temperatures than the eggs or young.
17. A benefit of heated effluents is the defouling of intake pipes accomplished by reversing the flow of water through the pipes.
18. Certain benefits, including open water winter fishing in otherwise ice covered areas, and a cold water fisheries downstream from deep reservoirs, can be derived from artificially induced temperature changes. The benefits of fish being attracted to heated water in the winter months may be negligible compared to fish mortalities that may result when the fish return to the cooler water; lethal temperatures may result from heated discharges in the summer months.


I. CHEMICAL REACTIONS

Introduction

All the impurities contained in a water result from intimate contact of the water with such impurities during which a portion are dissolved or suspended. The process of solution is a chemical reaction which proceeds as long as the water is in contact with a soluble substance or until equilibrium is reached (Camp, 1963). In general the solubility of solids and liquids may be considered a function of temperature, unless extreme pressure conditions are involved.

The solubility of nonreactive gases (gases that do not react with water to an appreciable extent) at equilibrium with the atmosphere is proportional to the partial pressure of the gas in the atmosphere and follows Henry's law, $C_s = K_s P$, where C_s is the saturation concentration of the gas in the water, P is the partial pressure of the gas phase and K_s is the proportionality constant called the coefficient of absorption. Water is saturated with a gas when the proportionality implied in Henry's law is fully established. Rising temperatures decrease the saturation value as do the salts of hard and brackish waters (Fair and Geyer, 1954). The solubility of reactive gases in water is modified because they ionize in and/or react with the water.

The establishment of equilibrium in a given chemical reaction implies that the reaction is reversible and that a point has been reached where a balance between the reactants exists. Mathematically, this relation is expressed by the mass action law



where capital letters refer to types of molecules or ions taking part in the reaction and lower case letters to the number of them. On the basis of thermodynamic principles the relation

$$\frac{(C)^p(D)^q}{(A)^m(B)^n} = k$$

is universally true, the parentheses indicating activities of the enclosed substances. The equilibrium constant k has a characteristic value for each reaction that is dependent only on temperature (Fair and Geyer, 1954).

The effects of temperature on equilibria are given by the Van't Hoff equation $\frac{d(\log_e K)}{dT} = \frac{\Delta H^\circ}{RT^2}$

where K is the equilibrium constant, T is the Kelvin temperature, R is the gas constant, ΔH° is the enthalpy change per gram-molecular weight for the reaction from left to right. Integration of the above equation between the limits T_1 and T_2 gives

$$\log_e \frac{K_2}{K_1} = \frac{\Delta H^\circ}{RT_1} - \frac{\Delta H^\circ}{RT_2} = \frac{\Delta H^\circ (T_2 - T_1)}{RT_1 T_2}$$

for constant ΔH^\ddagger and conversion to common logarithms

$$\text{Log} \frac{k_2}{k_1} = -\frac{\Delta H^\ddagger}{2.303R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right) = \frac{\Delta H^\ddagger}{4.576} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$

(Glasstone and Lewis, 1960).

The rates of most chemical reactions increase as the temperature is raised. A frequently used very approximate rule, enunciated by Van't Hoff, is that the rate doubles for each rise in temperature of 10°C (18°F).

Mathematically, the change in specific rate constant with temperature for any simple chemical reaction is given by the Arrhenius equation

$$\frac{d(\log_e k)}{dT} = \frac{E}{RT^2}$$

where k is the specific reaction rate constant, T is the Kelvin temperature, R is the gas constant (1.99 cal/degree C), E is a constant characteristic of the reaction and termed the activation energy.

Integration of the above equation between the limits T_1 and T_2 gives

$$\text{Log}_e \frac{k_2}{k_1} = \frac{E}{RT_1} - \frac{E}{RT_2} = \frac{E(T_2 - T_1)}{RT_1 T_2}$$

for constant E and conversion to common logarithms

$$\text{Log} \frac{k_2}{k_1} = \frac{-E}{2.303R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right) = \frac{E}{4.576} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$

(Glasstone and Lewis, 1960).

Two other methods for expressing the temperature dependence of a reaction rate are often encountered. They are:

$$\frac{k_2}{k_1} = \Theta^{T_2 - T_1}$$

whence

$$\Theta = \frac{k_2}{k_1} \text{ for } T_2 - T_1 \quad \text{and} \quad Q_{10} = \frac{k_2}{k_1} \text{ for } T_2$$

here T_2 and T_1 are measured in degrees centigrade (Fair and Geyer, 1954).

Dissolved Gases

1. Non-Reactive Gases

The gases that do not react to an appreciable extent with water but which occur in sufficient quantities to be determined by chemical analyses are oxygen, nitrogen, hydrogen and methane.


Oxygen

Since all living organisms are dependent on oxygen in one form or another to maintain the metabolic processes that produce energy for growth and reproduction, it is of great significance in the aquatic environment. The solubility of atmospheric oxygen (see Figure 1) in fresh waters ranges from 14.6 mg/l at 0°C (32°F) to 6.6 mg/l at 40°C (104°F) under 1 ATM of pressure (Anon., 1965). It is a poorly soluble gas and its solubility varies directly with the atmospheric pressure at any given temperature as expressed in Henry's law.

Oxygen solubility is affected also by the presence of dissolved salts as in hard and brackish waters. In brackish waters containing 5,000 mg/l of chloride, the solubility of atmospheric oxygen varies from 13.8 mg/l at 0°C (32°F) to 7.3 mg/l at 30°C (86°F) under 1 ATM of pressure (Anon., 1965).

In thermally stratified lakes, the hypolimnion may have a low oxygen content during summer stagnation because of its use in decomposition of organic matter. In addition, stratification may occur in streams when heated effluents are discharged. There are three recognized stream stratification forms: overflow, interflow, and underflow; the form being determined by the relationship between the density of the effluent and the density of the stream flow (Dysart and Krienkel, 1965). The principal source of dissolved oxygen is the air; oxygen

FIGURE 1 SOLUBILITY OF OXYGEN IN WATER
EXPOSED TO WATER-SATURATED AIR AT 760 mm Hg.



is also produced by chlorophyll-bearing algae and submerged aquatic plants through photosynthesis.

In natural waters deficient in dissolved oxygen, accelerated atmospheric aeration (reaeration) is evident. The rate of reaeration in each unit of time is proportional to the remaining degree of unsaturation of dissolved oxygen in the waters (Streeter, 1958). Temperature influences the rate of solution and affects the rate of diffusion of oxygen. As temperature increases the saturation capacity declines and the rate of diffusion increases (Velz and Gannon, 1960).

Nitrogen

The solubility of atmospheric nitrogen in water is about one-half that of oxygen. While the principal source of nitrogen is the air, denitrifying bacteria will release nitrogen to water (Allee et al., 1949). In a lake the waters of the hypolimnion become and remain supersaturated with nitrogen as they get warmer; rapid warming may cause nitrogen to escape as bubbles (Nordell, 1961).

Methane

Methane is not a permanent constituent of the earth's atmosphere (Camp, 1963). The primary source of methane in natural waters is anaerobic decomposition. Solubility of methane varies from 39.6 mg/l at 0°C (32°F) to 15.9 mg/l at 40°C (104°F) and 760 mm of mercury

(Nordell, 1961). Some ground waters may contain sufficient methane to constitute a fire and explosion hazard. Methane may persist even after aeration.

Hydrogen

The solubility of hydrogen varies from 1.93 mg/l at 0°C (32°F) to 1.48 mg/l at 40°C (104°F) in pure water in contact with the pure gas at 760 mm of mercury (Nordell, 1961). Hydrogen comprises less than 0.001% of the earth's atmosphere (Camp, 1963). It is produced in the water primarily from anaerobic decomposition of organic matter.

Dissolved Organic Substances



Organic substances in surface waters decay or decompose chiefly by bacterial action and exert a demand on the dissolved oxygen of such waters. This biochemical reaction is similar to an unimolecular chemical reaction, that is, the rate is approximately proportional to the remaining concentration of unoxidized organic matter. Thus the biochemical oxygen demand of a surface water is a measure of the concentration of decomposable organic matter (Camp, 1963). The reaction rate ordinarily expressed as k rises as the temperature of the water increases reaching a maximum at about 30°C (86°F), (Hoak, 1961). This increased rate results in a greater demand on the dissolved oxygen in the surface water. Theriault (1927) has shown that at 20°C (68°F) the biological oxidizability of polluted water increases by about 2% for each degree centigrade increase.

2. Reactive Gases

Gases that react with water are hydrogen sulfide, carbon dioxide, sulfur dioxide and ammonia. These either ionize in water or react with the water to produce ions.

Hydrogen Sulfide

The principal sources of hydrogen sulfide in natural waters are anaerobic decomposition of organic matter and the discharge of industrial wastes from oil refineries, leather tanneries, chemical plants and paper mills (Camp, 1963); the solubility of hydrogen sulfide ranges from 7,070 mg/l at 0°C (32°F) to 2,360 mg/l at 40°C (104°F) and 760 mm of mercury (Nordell, 1961). It ionizes in water in two steps--



An equilibrium relation is established for each step as follows:

$$1. \frac{[\text{H}^+][\text{HS}^-]}{[\text{H}_2\text{S}]}$$

$$2. \frac{[\text{H}^+][\text{S}^{=}] }{[\text{HS}^-]}$$

with equilibrium constants of 1.1×10^{-7} at 25°C (77°F) for step 1 and 1×10^{-14} at 25°C (77°F) for step 2. Equilibrium constants vary with temperature and corresponding values at 18°C (64.4°F) are 5.7×10^{-8} and 1.2×10^{-15} (Lange, 1961). As indicated by the comparative magnitude of the equilibrium constants for hydrogen sulfide at 25°C (77°F)

the equilibrium in step 1 lies to the side of the product, i.e. H^+ and HS^- , while the equilibrium in step 2 lies to the side of the reactant, i.e. HS^- . Because the equilibrium constants for steps 1 and 2 at $18^\circ C$ ($64.4^\circ F$) are less than the corresponding values at $25^\circ C$ ($77^\circ F$) it follows that reactions 1 and 2 decrease with a decrease in temperature (Prupton, 1951).

Sludge deposits in streams and estuaries produce hydrogen sulfide as they undergo anaerobic decomposition in which sulfates are reduced. Wheatland (1954) has indicated that the rate of formation of sulfide increases with temperature, doubling approximately for each $10^\circ C$ ($18^\circ F$) rise, and that reduction of sulfate to sulfide will occur at temperatures as low as $5^\circ C$ ($41^\circ F$); but even at $25^\circ C$ ($77^\circ F$) is inhibited by traces of dissolved oxygen. Hydrogen sulfide is oxidized in the presence of dissolved oxygen to water and free sulfur or to sulfate.

Carbon Dioxide


Free carbon dioxide is found in most surface waters and may range from 0 to 5 mg/l in rivers. Lake waters may contain from 0 to 2 mg/l at the surface with significant increases as the depth increases because of the processes of decay at or near the bottom (Allee, et al., 1949). Since the oxidation of organic matter furnishes carbon dioxide, much higher concentrations may be found in surface waters receiving organic wastes. Surface waters receiving acid mine drainage may show a high content of carbon dioxide; ground waters contain appreciable

amounts of free carbon dioxide, ranging from 1 mg/l to several hundred (Nordell, 1961).

The solubility of pure carbon dioxide in water ranges from 3,350 mg/l at 0°C (32°F) to 970 mg/l at 40°C (104°F) and 760 mm of mercury (Nordell, 1961). The average carbon dioxide content of the air varies from 0.035 percent in the country to 0.06 percent in the cities and the solubility of atmospheric carbon dioxide in water ranges from 1.0 mg/l at 0°C (32°F) and 0.03 percent to 2.0 mg/l at 0°C (32°F) and 0.06 percent. At 40°C (104°F) the solubility is 0.3 mg/l and 0.6 mg/l respectively at 760 mm of mercury (Nordell, 1961). These values indicate that the carbon dioxide contributed to natural surface waters and ground waters from the atmosphere is negligible compared with that from decaying organic matter.

Carbon dioxide reacts with water to form a weakly dissociated acid-carbonic acid, $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3$

Carbonic acid is a dibasic acid ionizing in two steps:



An equilibrium relation is established for each step as follows:

$$1. \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$$


$$2. \frac{[\text{H}^+][\text{CO}_3^{--}]}{[\text{HCO}_3^-]}$$

Equilibrium constant values for steps 1 and 2 respectively, are 4.31×10^{-7} and 5.6×10^{-11} at 25°C (77°F) (Lange, 1961). At this point carbon dioxide establishes an equilibrium relation with the mineral content of the water.

Ammonia

Ammonia is an intermediate product in the bacterial decomposition of nitrogenous organic matter and may be discharged to natural waters as a waste product of industry. The presence of free ammonia in natural waters is indicative of recent organic pollution, since the atmosphere is substantially free of this substance (Camp, 1963).

The gas reacts with water to produce ammonium hydroxide (often termed aqueous ammonia); the ammonium hydroxide in turn ionizes to produce ammonium and hydroxal ions.



An equilibrium relation is established as follows:

$$\frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]}$$



With ionization constant values of 1.8×10^{-5} at 25°C (77°F) and 2.0×10^{-5} at 40°C (104°F) (Lange, 1961). The equilibrium is shifted toward the product side, i.e. NH_4^+ ; OH^- at higher temperatures. Because the hydroxyl ion is a product of the ionization of ammonium hydroxide, the degree of ionization can be related to the hydrogen ion concentration or pH. Camp (1963) indicates that 99.99% of the ammonia in

dilute solutions at 25°C (77°F) is in the form of ammonium ion at a pH of 5 while at pH 11 only 1.78% is in this form.

Sulfur Dioxide

Sulfur dioxide is formed in one of three ways: (1) as an intermediate product in the oxidation of hydrogen sulfide under aerobic conditions, (2) in the reduction of sulfate under anaerobic conditions and (3) in the combination of elemental sulfur and oxygen. This is part of the sulfur cycle taking place in many natural waters (Sawyer, 1960). Combustion fumes from industrial operations may contribute to the sulfur dioxide content of natural waters.

Sulfur dioxide reacts with water to form sulfurous acid; the acid then ionizing in two steps--



and equilibrium relation is established as follows:

$$1. \frac{[\text{H}^+][\text{HSO}_3^-]}{[\text{H}_2\text{SO}_3]}$$

$$2. \frac{[\text{H}^+][\text{SO}_3^{--}]}{[\text{HSO}_3^-]}$$

With ionization constant values of 1.72×10^{-2} and 6.24×10^{-8} at 25°C (77°F) for steps 1 and 2, respectively (Lange, 1961). In the presence of dissolved oxygen, sulfurous acid and the hydrogen sulfite ion are readily oxidized. The solubility of sulfur dioxide markedly decreases with increasing temperatures (Camp, 1963).

Dissolved Minerals

About 99% of the dissolved mineral matter found in natural waters embrace only 10 elements, namely; hydrogen, oxygen, sodium, potassium magnesium, calcium, silicon, sulfur, carbon and chlorine. These occur as ions, radicals or molecules. At ordinary temperatures of natural waters some complexing of major dissolved species may occur but is limited to the formation of ion pairs [e.g., $\text{Na}(\text{CO}_3)_2^-$, $\text{Na}_2\text{CO}_3^\circ$] (Garrels and Christ, 1965). Helgeson (1964) in a study of the effects of elevated temperatures on the dissociation of complex ions in solution indicated that little or no changes occur unless there is an appreciable change in the density of the solution; with increasing temperature and decreasing density complexing is expected to increase.

At present only enough chemical information is available to permit calculation of the inter-actions that take place among the major dissolved species at earth surface temperatures in media as concentrated as sea water. The results of such calculations indicate that more than 30% of the sulfate and bicarbonate are tied up as ion pairs with cations, whereas 90% of the total carbonate is complexed. One-hundred percent of the chloride is present in the ionic form. Changes in temperature, pressure and composition of the water will modify this distribution; however, variations in temperature and pressure to which ocean waters are subjected will produce little change in distribution.

The effects of small variations in temperature [$5^{\circ}\text{C}(41^{\circ}\text{F})$ to $40^{\circ}\text{C}(104^{\circ}\text{F})$] on equilibria and reaction rates involving minor constituents in natural waters can not be determined at this time because of a lack of information on equilibrium constants, enthalpy changes and activation energies (Garrels and Christ, 1965).

Summary

1. Chemical reaction rates vary with temperature, generally increasing as the temperature is increased. The change in the specific rate constant is given by the Arrhenius equation.
2. The solubility of gases in water varies with temperature. Dissolved oxygen content of a surface water is decreased by the decay or decomposition of dissolved organic substances; the decay rate increases as the temperature of the water increases reaching a maximum at about 30°C (86°F).

REFERENCES CITED

- Allee, W. C., et al., 1949. Principles of animal ecology. W. B. Saunders Co., Philadelphia, 837 pp.
- Anon., 1965. Standard methods for the examination of water and wastewater. Twelfth Edition, American Publ. Health Assn., Inc., New York, 769 pp.
- Camp, T. R., 1963. Water and its impurities. Reinhold Publishing Corp., New York, 355 pp.
- Dysart, B. C., and P. A. Krenkel, 1965. The effects of heat on water quality. Proc. 20th Ind. Waste Conf., Purdue Univ., pp. 18-39.
- Fair, G. M. and J. C. Geyer, 1954. Water supply and wastewater disposal. John Wiley & Sons, Inc., New York, 973 pp.
- Garrels, R. M. and C. L. Christ, 1965. Solutions, minerals, and equilibria. Harper & Row, New York, 450 pp.
- Glasstone, S. and D. Lewis, 1960. Elements of physical chemistry. D. Van Nostrand Co., New York, 758 pp.
- Helgeson, H. C., 1964. Complexing and hydrothermal ore deposition. Pergamon, New York, 128 pp.
- Hoak, R. D., 1961. The thermal pollution problem. Jour. Water Poll. Control Fed., 33(12): 1267-1276.
- Hutchinson, G. E., 1957. A treatise on limnology. John Wiley & Sons, Inc. New York, I: 1015 pp.
- Lange, N. A., 1961. Handbook of chemistry, tenth edition. McGraw-Hill Book Co., New York, 1969 pp.
- Nordell, E., 1961. Water treatment for industrial and other uses. Reinhold Publishing Corp., New York, 598 pp.
- Prutton, C. F. and S. H. Marion, 1951. Fundamental principles of physical chemistry. The MacMillan Co., New York, 803 pp.

Sawyer, C. N., 1960. Chemistry for sanitary engineers. McGraw-Hill Book Co., New York, 367 pp.

Streeter, H. W., 1958. The oxygen sag and dissolved oxygen relationships in streams. Oxygen Relationships in Streams, U. S. Dept. of Health, Education, and Welfare, Robert A. Taft Sanitary Engineering Center, Cincinnati, 25-30.

Theriault, E. J., 1927. The oxygen demand of polluted water. U. S. Public Health Service Bull. 173.

Velz, C. J. and J. J. Gannon, 1960. Forecasting heat loss in ponds and streams. Jour. Water Poll. Control Fed., 32(4): 392-417.

Wheatland, A. B., 1954. Factors affecting the formation and oxidation of sulfides in a polluted estuary. Jour. of Hygiene, Cambridge Univ. Press, 194-210.

II. BACTERIA

Introduction

Temperature changes in the aquatic environment affect ecological relationships among the biota, processes of natural purification, and growth and survival of microorganisms. There are a wide variety of microorganisms found in the aquatic environment. The numbers and species in the population vary depending on whether they are in ground water, lakes, or streams. Unpolluted water bodies have low concentrations of microorganisms. The microbial content of natural waters is approximately proportional to the amount of organic matter present. Unpolluted waters usually have a greater number of species in proportion to their total population; conversely polluted waters usually have a greater total population in proportion to the number of bacterial species in their environment.

Many stream bacteria come from the air and soil. Bacteria in the air are aerosols or suspended on dust particles that settle or fall with precipitation. Soil flora in water are due to precipitation and seeping ground water that becomes surface run-off when entering or forming a stream. Many of the microorganisms that are native to natural waters are especially adapted to the stream environment, and some are difficult or now impossible to grow on culture media.

Interest in the microbiology of water centers on the transmission of disease via the water route. Polluted waters have high concentrations of microorganisms from municipal or industrial waste waters. When a stream is polluted with the excreta of warm blooded animals, it is most likely to contain enteric pathogens. Indicator organisms (coliforms) are

used to measure bacterial concentrations and indicate the potential presence of pathogens from sewage.

Industrial wastes add organic and inorganic materials to water and in some cases large numbers of bacteria, but, in general, they do not contain pathogenic organisms. Industrial wastes may be growth stimulating or toxic to bacteria (Heukelekian, 1953).

Water, an essential for all forms of life, can serve as a medium for growth and reproduction for many microorganisms. Environmental factors affecting the growth of microorganisms are chemical, physical, and nutritional. Although these factors are interdependent, the physical factor of temperature is one of the most important.

The relations of temperature to the growth of microorganisms are complex. Some hardy bacteria grow in a wide temperature range; other fragile bacteria grow in a narrow temperature range. For each organism there is a minimum, the lowest temperature at which growth can occur; an optimum, the temperature most favorable for growth; and a maximum, the highest temperature at which growth and multiplication can occur.

Bacteria may be classified according to their temperature requirements for growth. Organisms having optimum growth temperatures under 20°C (68°F) are grouped as psychrophiles or cold-loving; these occur in the soil and cold waters of the north. Thermophiles, heat-loving organisms having optimum temperatures of 55° to 65°C (131 to 149°F) are found in soil, decaying organic matter, hot spring water and near the discharge points of hot water effluents. They are of little importance in stream ecology. The majority of bacteria are called mesophiles; they are the intermediate group, having optimum temperatures in the range between the extremes. Many of the organisms found in natural waters and

soils are saprophytes (organisms that live on decaying organic matter); they have optimum temperatures of 22° to 28°C (70 to 82°F) and belong to the mesophilic group.

The parasitic bacteria have optimum temperatures of 37°C (98.6°F) and include those microorganisms pathogenic to man. Temperature changes greatly affect the rate of activity of these organisms.

The effect of temperature on the species of organisms cannot always be considered separately from the other environmental factors. Some species can be more abundant in the winter, others in the summer, when their environmental conditions are varied (Burrows, 1959).

Microbiotic Cycles

The interrelationships among aquatic biota are of primary importance to the aquatic environment (Ingram, Mackenthun, and Bartsch, 1966). In unpolluted waters, autotrophic algae and chlorophyll bearing bacteria initiate microbiotic cycles (Silvey and Roach, 1964). The metabolites and decomposition products of the organisms provide nutrients for use by gram-negative heterotrophic bacteria. The principal gram-negative organisms are Alcaligenes, Aerobacter, and Pseudomonas. In waters polluted with sewage, the Escherichia also increase, as do associated types. The new algal growth following the gram-negative bacilli may be either diatoms or blue-green algae. In cool waters the diatoms will generally prevail and be followed in turn by the gram-positive heterotrophic bacilli. In warmer waters, the gram-negative bacilli remain and grow in coexistence with blue-green algae; then the actinomycetes grow and develop an antibiotic effect, reducing the gram-negative population. The gram-positive spore-forming bacilli follow the actinomycetes (Figure 2).




Fig. 2. Annual Cycles of Gram-Positive and Gram-Negative Heterotrophic Bacilli and Their Relationship to Blue-Green Algae and Aquatic Actinomycetes

The dotted curve is for blue-green algae, indicated in 100's of areal standard units on Scale 1 at the left; the solid curve is for gram-positive heterotrophs; the dashed curve, gram-negative heterotrophs; and the dot-and-dash curve, actinomycetes. Colony counts of heterotrophs are indicated in 1,000's by Scale 1; absolute numbers of bacilli by Scale 2; and actinomycete plate counts in 1,000's by Scale 3. Actinomycetes were isolated on M_1B_2 agar. Gram-positive heterotrophs were grown on vitamin-enriched Emerson's agar, and gram-negative heterotrophs on eosin methylene blue agar.

(From Silvey and Roach, 1954, p. 54)

These various cycles may impinge upon each other when the stream or reservoir is highly polluted.

Self-Purification

When organic wastes are discharged into a receiving water a complex chain of physical, chemical, biochemical and biological activities are started which result in decomposition and degradation of the wastes. This complex process is self-purification, the details of which are still unknown (Heukelekian, 1953).

The most significant function in natural purification is the decomposition of organic matter by the microbial flora. The saprophytic organisms are the most active in this biochemical process; they have optimum temperatures that are near the ambient temperature of many streams during the summer months. In a stream polluted with sewage, the pathogenic and indicator organisms are also present and perform a minor role in the self-purification process. The temperature of the stream water even during the summer is below the optimum for pollution associated bacteria. Increasing the water temperature increases the bacterial multiplication rate when the environment is favorable and the food supply is abundant. Increasing the water temperature within the growth range of the bacteria causes a more rapid die-off when the food supply is limited. The decrease of bacterial numbers is higher during the summer than during the winter (Figure 3).

Unpolluted streams contain dissolved oxygen near saturation levels. When organic wastes are discharged into the stream, the biochemical process is aerobic - reducing the dissolved oxygen by oxidation and dilution. When the reaeration rate of the stream is low, the oxygen may be depleted by the bacterial metabolism of the increasing population.




Figure 3. From Huekelekian, 1953, p. 27

As the available oxygen diminishes, the aerobic organisms die-off rapidly, sharply decreasing the natural purification process of assimilating the organic waste load. When this occurs there is a shift in the flora in the stream to the facultative anaerobic organisms. The ultimate result of anaerobic decomposition may be the same as that of the aerobic, but it is very slow and less desirable (Fair, Geyer, and Morris, 1958).

As the temperature increases, the dissolved oxygen solubility decreases. When a warm water discharge is near a sewage treatment plant outfall, self-purification can be very effective if the organic waste load is not excessive.

The impounding of water may improve the water quality by reducing sediments, color, bacteria and temperature. Ingols (1957) found also that the pattern of reservoir discharge permitted slime development downstream during low flow and scouring of the slime during high flows. "Out of phase" dilution of the receiving stream could promote or retard stream self-purification (Berger, 1961). Berger as well as Renn (1957) agree that the rate of stream reaeration increased at higher temperatures.

Growth and Survival

Chambers and Clarke (1966) state: "Many bacteria reproduce in water: among the genera that will grow in water of unquestioned potable quality are: (1) Pseudomonas, (2) Xanthomonas, (3) Achromobacter, (4) Escherichia, (5) Aerobacter, (6) Streptococcus, (7) Desulfovibrio, and (8) Crenothrix."

Renn (1957) points out that elevating the stream temperature can be favorable for those bacteria that can multiply in water by inducing

the recurring cycles of life and death more rapidly. However, enteric pathogens have highly selective requirements. They cannot multiply or survive well in natural water, so they die-off more rapidly.

Because higher temperatures in a stream polluted with sewage generally result in increased bacterial numbers, low temperatures are not conducive to rapid growth. Stream temperatures of 1° to 8°C (33.8 - 46.4°F) may suppress growth and multiplication, and act as a preservative as in the storage of samples for bacterial analyses. Freezing of water can result in reduced microbial populations by killing off a majority of the microorganisms. Streams that have high organic waste loads and low temperatures tend to develop slime organisms, generally Sphaerotilus.

A study of the Columbia River in Oregon showed Sphaerotilus growths to be maximum at water temperatures of 10° to 15°C (50° - 59°F). Growth ceased when temperatures dropped below 4°C (39.2°F) and resumed when temperatures increased above 4°C (39.2°F). Infestations of Sphaerotilus may occur at temperatures below 10°C (50°F) if the growing period is sufficiently long (Amberg and Cormack, 1960). Beds of Sphaerotilus slime may extend farther downstream from a waste outfall in the winter than in the summer when warmer temperatures seem to inhibit the efficiency of food conversion by the organism (Dondero, 1961).

Clark et al. (1964) assessed the value of bacterial indicators of pollutions as indicators of viral pollution by studying the relative survival of the organisms in water. They observed that the lower the temperature the longer the survival for both bacteria and viruses. The enteric bacteria had survival times in proportion to the degree of pollution, the greater the pollution the longer the survival time. The

increased quantity of nutrients present in the more polluted water may account for the longer survival time of the bacteria. The viruses studied survived longer in the "clean" Little Miami River water and in the grossly polluted raw sewage than in the moderately polluted Ohio River water. They also point out the difficulty in generalizing on comparative survival times because the different genera of organisms may have different survival times in the different stream environments (Table 1).

Summary

The temperature of stream water, even during the summer, is below the optimum for pollution-associated bacteria. Increasing the water temperature increases the bacterial multiplication rate when the environment is favorable and the food supply is abundant. Increasing the water temperature within the growth range of the bacteria causes a more rapid die-off when the food supply is limited.

TABLE 1

AVERAGE TIME IN DAYS FOR 99.9 PERCENT REDUCTION IN ORIGINAL
TITER OF INDICATED MICROORGANISMS AT THREE TEMPERATURES

(From Clarke, Berg, Kabler and Chang, p. 526, 1964)

Microorganisms	Little Miami River			Ohio River			Sewage		
	28°C	20°C	4°C	28°C	20°C	4°C	28°C	20°C	4°C
<u>Poliovirus I</u>	17	20	27	11	13	19	17	23	110
<u>ECHO 7</u>	12	16	26	5	7	15	28	41	130
<u>ECHO 12</u>	5	12	33	3	5	19	20	32	60
<u>Coxsackie A9</u>	< 8	< 8	10	5	8	20	6	No Data	12
<u>A. aerogenes</u>	6	8	15	15	18	44	10	21	56
<u>E. coli</u>	6	7	10	5	5	11	12	20	48
<u>S. fecalis</u>	6	8	17	9	18	57	14	26	48

REFERENCES CITED

- Amberg, H. R. and J. F. Cormack, 1960. Factors affecting slime growth in the lower Columbia River and evaluation of some possible control measures. Pulp Paper Mag. Can., 61:T-70 to T-80.
- Berger, B. B., 1961. Does production of power pollute our rivers? Power Engineering, March, p 60-61.
- Burrows, W., 1959. Textbook of microbiology. W. B. Saunders Co., Philadelphia, Pa.
- Chambers, C. and N. Clarke, 1966. Control of bacteria in nondomestic water supplies. Advances in Applied Microbiology. Academic Press Inc., N. Y., 8:105-143.
- Clarke, N., G. Berg, P. Kabler, and S. Chang, 1964. Human enteric viruses in water: source, survival and removability. Reprint from International Conference on Water Pollution Research. London, 1962. Pergamon Press - Oxford, London, N. Y. and Paris, pp. 523-536.
- Donders, N. C., 1961. Sphaerotilus, its nature and significance. In: Advances in Applied Microbiology, 3:77-107.
- Fair, G. M., J. C. Geyer and J. C. Morris, 1958. Water supply and waste-water disposal. John Wiley & Sons, Inc., New York.
- Heukelekian, H., 1953. Stream pollution and self-purification. In: industrial wastes their disposal and treatment. (W. Rudolfs. Ed.) Reinhold Pub. Corp., New York, pp 8-30.
- Ingols, R. S., 1957. Pollutational effects of hydraulic power generation stream pollution. Sewage and Industrial Wastes, 29(3):292-297.
- Ingram, W. M., K. M. Mackenthun, and A. F. Bartsch, 1966. Biological field investigative data for water pollution surveys. U. S. Department of the Interior, Federal Water Pollution Control Administration, WP-13, 139 pp.
- Renn, C. E., 1957. Warm-water effects on municipal supplies. Jour. American Water Works Assoc., 49,405-412.
- Silvey, J. K. G., A. W. Reach, 1964. Studies on microbiotic cycles in surface waters. Jour. American Water Works Assoc., 56(1):60-72.

III. FRESHWATER FISHES

Introduction

Changes in fish populations can result from the many types of artificial cooling and heating of natural waters. These changes result from the discharge of condensed water from steam-electric generating plants, distillery effluents, and irrigation waters. Stream temperatures are raised also by the removal of stream bank trees and other vegetation. Water temperatures are often elevated in excess of the air temperature by absorption of heat by the stream bed. Yet another type of thermal pollution results from the discharge of cold-water from stratified impoundments; this water may provide an ideal habitat for trout and other cold-water fish when sufficient dissolved oxygen is present.

General Effects

The effects of temperature on fish are acute because fish do not possess an efficient method to compensate their internal temperature against a temperature change in the water in which they are immersed. If a more favorable temperature is available, fish do have the ability to seek it out.

Chemical reactions are accelerated within the body cells with elevated temperatures. Prosser (1955) discusses four possible death mechanisms, although he does not attach specific temperature values to various death processes. These are: an enzyme inactivity caused by the acceleration of

the enzyme reaction to such a state that it is no longer effective; coagulation of cell proteins; melting of cell fats; reduction in the permeability of cell membranes. Cells may also be killed by toxic action of the products of metabolism and, incomplete metabolism accumulating in the cells (Ellis, 1947).

According to Brett (1960), temperature acts in a variety of different ways; it can be lethal, cause a reduction of activity, and limit reproduction. The slow rate of acclimation appears to result in greater mortalities from cold despite the ability of fish to withstand lower temperatures.

Sudden Temperature Changes

The effect of thermal shock on fish can be more harmful than continued exposure to a higher temperature (Cairns, 1956). In studies with rainbow trout, Threinen (1958) found that death would result from an instantaneous or rapid increase (shock) of 11.1°C (20°F) above an acclimation temperature of 12.2°C (54°F), however, a similar increase of 8.4°C (15°F) could be tolerated from a temperature of 10.6°C (51°F). A rise of the acclimation temperature from 12.2 to 18.4°C (54 - 65°F) during a 24-hour period permitted trout to withstand a temperature of 23.4°C (74°F) with only minor distress for short periods.

Fish having the ability to adapt to higher temperatures faster and over a larger gradient often are attracted to artificially heated water without a resultant mortality. However, mortality often results when these fish

return to cold water. Agersborg (1930) found fish dying when they attempted to return from heated water (26.1°C - 79°F) to the colder stream (0°C - 32°F); death occurred even when fish moved into water that was 5.6°C (10°F) cooler. Falkner and Houston (1966) found that the mean erythrocytic (red blood cell) volume underwent a transient decrease while total blood iron (and presumably haemoglobin and mean erythrocytic iron content) fell slightly after goldfish which had been acclimated to 20°C (68°F) were subjected to an abrupt increase of 10°C (18°F). Heinicke and Houston (1965) concluded that while thermal shock induces initial deviations in iono- and osmoregulatory ability the goldfish can compensate for these changes during the acclimation period through respiratory activities, and restore its original ionic status.

A rise in temperature from 10°C to 20°C (50 - 68°F) reduced resistance to a decrease in oxygen in perch, roach, and mirrorcarp. In rainbow trout the resistance was lowered considerably between a rise in temperature from 10°C to 16°C (50 - 60.8°F) (Downing and Merkens, 1957).

Acclimation

The importance of acclimation temperatures has long been known to fish hatchery personnel and physiologists working on lethal temperatures. Much of the work on lethal temperatures is of little value because holding temperatures and durations are not given. Springtime mortalities often result from fish being subjected to warmer water temperature after acclimation to cold winter temperatures.

Doudoroff, in Brown (1957), discussing the work of Fry, et al., (1946) and Fry (1947) concludes that fish could stand brief exposures to considerably higher temperatures without showing distress when they had been acclimated to the maximum possible temperature. However, the fish suffered mortality when they had been acclimated to low temperatures. Similarly, Doudoroff in discussing the work of Hart (1952) noted considerable geographic, seasonal, and other variations of the resistance to heat of some species of fish acclimated to the same temperature.

Doudoroff in discussing the rate of acclimation summarized the work of many workers and concluded that the increased heat resistance (the ability to withstand increased water temperatures) is acquired usually at a very fast rate in the high temperature range from 26°C to 30°C (78.8-86.0°F) although there may be a latent period of one or longer days in which virtually no change takes place in the upper lethal temperatures. Most of the resulting increase was achieved in a period of one to three days. There was little or no loss of resistance in the first three days. Thus, if a fish has acquired a higher heat resistance it will not be lost rapidly on subsequent exposures to low temperatures.

Jones (1964) in discussing the work of Sumner and Wells noted that the tolerance to high temperatures once acquired may persist for considerable periods after return of the fish to the acclimation temperature. The time

of acclimation need not be continuous. An intermittent exposure to a different temperature for sufficient hours per day can produce the same acclimation temperature as a continuous exposure.

The acclimation of fish is important in determining the maximum environmental temperature in which fish can survive. Jones (1964) discusses how the resistance time shortens with a progressive rise in temperature until the fish succumbs to an ultimate lethal temperature. As the acclimation temperature rises the thermal death point rises, but it rises at a slower rate. Accordingly, experiments on roach show that for every 3-degrees rise in acclimation temperature the thermal death point rises only 1-degree C (1.8°F).

Maximum Temperatures

Maximum temperatures have been determined for numerous species of fish (Table 2). These temperatures are important in determining the absolute temperature at which a fish can survive, but they are often higher than the maximum temperature at which a population can survive.

Alabaster (1962) found that heated effluents, by virtue of their high temperatures only, may be lethal to caged trout and coarse fish acclimated to normal river temperatures during the summer and may also occasionally kill free-living fish which are near effluent outfalls when temperatures increase rapidly. Small free-living fish are principally affected, large fish apparently are able to swim away to safety. He concludes further that where the water temperature of the whole river is above normal because of mixing with continuous discharge of heated effluent,

TABLE 2

TOLERANCE LIMITS FOR CERTAIN FISHES

Values are LD₅₀ temperature tolerance limits, i.e., water temperatures survived by 50 percent of the test animals. Counts were made by observing or estimating the number killed during exposure, or within a reasonable time thereafter in which it could be safely assumed that all deaths were attributable to the temperature effects.

(This Table Taken in Part From Anon., 1962)

Fish	Acclimated to		Lower Limit		Hr	Upper limit		Hr
	°C	(°F)	°C	(°F)		°C	(°F)	
Bass, largemouth (<u>Micropterus salmoides</u> <u>floridanus</u>)	20.0°C	(68.0°F)	5.0°C	(41.0°F)	24	32.0°C	(89.6°F)	72
	30.0°C	(86.0°F)	11.0°C	(51.8°F)	24	24.0°C	(93.2°F)	72
Bluegill (<u>Lepomis</u> <u>macrochirus macrochirus</u>)	10.0°C	(50.0°F)				29.0°C	(82.4°F)	24
	30.0°C	(86.0°F)				36.0°C	(96.9°F)	24
Bluegill (<u>L. macrochirus</u> <u>purpureus</u>)	15.0°C	(59.0°F)	3.0°C	(37.4°F)	24	31.0°C	(87.8°F)	60
	30.0°C	(86.0°F)	11.0°C	(51.8°F)	24	34.0°C	(93.2°F)	60
Bullhead (<u>Ameiurus n.</u> <u>nebulosus</u> , A. n. <u>narmoratus</u>)	20.0°C	(68.0°F)	1.0°C	(33.8°F)	24	32.0°C	(89.6°F)	96
	30.0°C	(86.0°F)	7.0°C	(44.6°F)	24	35.0°C	(95.0°F)	96
Catfish, channel (<u>Ictalurus lacustris</u> <u>lacustris</u> , I. l. <u>punctatus</u>)	15.0°C	(59.0°F)	0.0°C	(32.0°F)	24	30.0°C	(86.0°F)	24
	25.0°C	(77.0°F)	6.0°C	(42.8°F)	24	34.0°C	(93.2°F)	24
Chub, creek (<u>Semotilus</u> <u>a. atromaculatus</u>)	5.0°C	(41.0°F)				25.0°C	(77.0°F)	96
	25.0°C	(77.0°F)				32.0°C	(89.6°F)	96
Dace, blacknose (<u>Rhinich-</u> <u>thys a. atratulus</u> , R. a. <u>meleagris</u>)	5.0°C	(41.0°F)				30.0°C	(80.6°F)	340
	25.0°C	(77.0°F)	5.0°C	(41.0°F)	24	29.0°C	(84.2°F)	340
Goldfish (<u>Carassius</u> <u>auratus</u>)	2.0°C	(35.6°F)				28.0°C	(82.4°F)	14
	17.0°C	(62.6°F)	0.0°C	(32.0°F)	14	34.0°C	(93.2°F)	14
	24.0°C	(75.2°F)	5.0°C	(41.0°F)	14	36.0°C	(96.8°F)	14
Greenfish (<u>Girella</u> <u>nigricans</u>)	37.0°C	(98.6°F)	15.0°C	(59.0°F)	14	42.0°C	(107.6°F)	14
	12.0°C	(53.6°F)	5.0°C	(41.0°F)	120	30.0°C	(86.0°F)	120
	18.0°C	(64.4°F)	13.0°C	(55.4°F)	72	31.0°C	(87.8°F)	120
Killifish (<u>Fundulus</u> <u>heteroclitus</u>)	14.0°C	(57.2°F)	1.0°C	(33.8°F)	48	32.0°C	(89.6°F)	
	20.0°C	(68.0°F)	2.0°C	(35.6°F)	48	34.0°C	(93.2°F)	
Minnow, fathead (<u>Pimephales promelas</u>)	20.0°C	(68.0°F)	2.0°C	(35.6°F)	24	32.0°C	(89.6°F)	133
	30.0°C	(96.0°F)	11.0°C	(51.8°F)	24	33.0°C	(91.4°F)	133
Minnow, blunt-nose (<u>Hyborhynchus notatus</u>)	15.0°C	(59.0°F)	1.0°C	(33.8°F)	24	31.0°C	(87.8°F)	133
	25.0°C	(77.0°F)	8.0°C	(46.4°F)	24	33.0°C	(91.4°F)	133

Fish	Acclimated to		Lower Limit		Hr	Upper Limit		Hr
	°C	(°F)	°C	(°F)		°C	(°F)	
Mosquito fish (<u>Grambusia affinis</u> <u>affinis</u> , G.a. <u>holbroki</u>)	15.0°C 35.0°C	(59.0°F) (95.0°F)	2.0°C 15.0°C	(35.6°F) (59.0°F)	24 24	35.0°C 37.0°C	(95.0°F) (98.6°F)	-66 .66
Perch (<u>Perca flavescens</u>)	5.0°C	(41.0°F)				21.0°C	(69.8°F)	96
Winter	25.0°C	(77.0°F)	4.0°C	(39.2°F)	24	30.0°C	(86.0°F)	96
Summer	25.0°C	(77.0°F)	9.0°C	(48.2°F)	24	32.0°C	(89.6°F)	96
Shad, gizzard (<u>Dorosoma cepedianum</u>)	25.0°C 35.0°C	(77.0°F) (95.0°F)	11.0°C 20.0°C	(51.8°F) (68.0°F)	24 24	34.0°C 37.0°C	(93.2°F) (98.6°F)	48 48
Shiner, common (<u>Notropis cornutus</u> <u>frontalis</u>)	5.0°C 25.0°C 30.0°C	(41.0°F) (77.0°F) (86.0°F)	4.0°C 8.0°C	(39.2°F) (46.4°F)	24 24	27.0°C 31.0°C 31.0°C	(80.6°F) (87.8°F) (87.8°F)	133 133 133
Shiner, common (<u>Notropis cornutus</u> <u>chrysocephalus</u>)	25.0°C 30.0°C	(77.0°F) (86.0°F)				32.0°C 34.0°C	(89.6°F) (93.2°F)	133 133
Shiner, lake (N. <u>atherinoides</u>)	5.0°C 15.0°C 25.0°C	(41.0°F) (59.0°F) (77.0°F)	2.0°C 8.0°C	(35.6°F) (46.4°F)	24 24	23.0°C 29.0°C 31.0°C	(73.4°F) (84.2°F) (87.8°F)	133 133 133
Shiner, golden (<u>Notemigonus</u> c. <u>crysoleucas</u> , N.c. <u>auratus</u>)	20.0°C 30.0°C	(68.0°F) (86.0°F)	8.0°C 11.0°C	(46.4°F) (51.8°F)	24 24	32.0°C 35.0°C	(89.6°F) (95.0°F)	66 66
Sucker, common (<u>Catostomus commersoni</u>)	15.0°C 25.0°C	(59.0°F) (77.0°F)	5.0°C	(41.0°F)	24	29.0°C	(84.2°F)	133
Sunfish (<u>Lepomis gibbosus</u>)	10.0°C 30.0°C	(50.0°F) (86.0°F)				28.0°C 24.0°C	(82.4°F) (75.2°F)	24 24
Trout, brook (<u>Salvelinus fontinalis</u>)	3.0°C 20.0°C 25.0°C	(37.4°F) (68.0°F) (77.0°F)				23.0°C 25.0°C 25.0°C	(73.4°F) (77.0°F) (77.0°F)	133 133 133

coarse fish populations may be reduced locally when the mean daily temperature reaches 30°C (86°F) and increased when the water is not warmed to more than 26°C (78.8°F).

Wells (1914) concluded that the resistance of fish to temperature varies with species and size of fish. There is no definite maximum temperature for a given species of fish; it varies with the fish's rate of heating, size, and physiological condition.

A temperature need not kill the fish directly for it to be lethal. Brook trout were found to be comparatively slow in catching minnows at 17.2°C (63°F) and virtually incapable of catching minnows at 21°C (70°F). This resulted in the trout virtually starving to death (Anon., 1962).

As a maximum temperature for cold water fishes the Pennsylvania Department of Health recommends*: that no wastes or waters shall be added from any source having temperatures in excess of that of the receiving waters except that during the period October through May, when stream temperatures are below 14.5°C (58°F) the temperature of wastes discharged to the streams shall not exceed 14.5°C (58°F). To allow for the normal production of aquatic life in warm water lakes and streams it is recommended that water temperatures resulting from thermal discharge shall not exceed

* Anon., 1962. Heated discharges . . . their effect on streams. Rep. by the Advisory Committee for the control of stream temperatures to the Pa. Sanitary Water Board. Pa. Dept. Health, Harrisburg, Publ. No. 3, 108 pp.

30°C (93°F) exclusive of the required mixing zone and in no case shall this peak temperature prevail for more than eight hours in any 24-hour period.

Preferred Temperature

It is generally acknowledged that fish can live for short periods of time in higher than normal temperatures, but at these temperatures fish cannot perpetuate their populations. Fish are extremely sensitive to temperature and seek out the temperature that is best for their survival. The temperature that fish seek out is termed "preferred temperature;" these are listed for several species of fish in Tables 3 and 4 and Figures 4 and 5.

Windermere char eggs hatched in 45 days at 10°C (50°F), and in 95 days at 4°C (39.2°F) (Swift, 1965). Mortality to some extent occurred at 8°C (46.4°F), with total mortality occurring at 12°C (53.6°F).

Ferguson (1958) concluded that the level of thermal acclimation influences the range of temperature preferred. In general, the preferred temperature is considerably higher than the acclimation temperature at lower thermal acclimations, but this difference decreases up to the final preferred temperature where both coincide. A final preferred temperature and the relation between acclimation and preferred temperature is characteristic for the species.

Tarzwel (1957) concluded that while temperatures higher than the optimum, and high temperatures of short duration, 23.9 to 27.8°C (75 to 82°F),

TABLE 3

THE FINAL TEMPERATURE PREFERENDA FOR VARIOUS SPECIES OF FISH
AS DETERMINED BY LABORATORY EXPERIMENTS

Young of the Year or Yearling Fish Were Used, Except as Noted.
(This Table Taken in Part From Ferguson, 1958)

Species	Final Preferendum	Authority
Bluegill (<u>Lepomis macrochirus</u>)	32.3°C (90.1°F)	Fry and Pearson (MS, 1952)
Bass, Largemouth (<u>Micropterus salmoides</u>)	30.0-32.0°C (86-89.6°F)	Fry (MS, 1950)
Carp (<u>Cyprinus carpio</u>)	32.0°C (89.6°F)	Pitt, Garside and Hepburn (1956)
Pumpkinseed (<u>Lepomis gibbosus</u>)	31.5°C (88.7°F)	Anderson (MS, 1951)
Goldfish (<u>Carassius auratus</u>)	28.1°C (78.8°F)	Fry 1947
Bass, Smallmouth (<u>Micropterus dolomieu</u>)	28.0°C (82.4°F)	Fry (MS, 1950)
Grass Pickerel (<u>Esox vermiculatus</u>)	26.6°C (78.8°F)	Berst and Lapworth (MS, 1950)
Yellow Perch (<u>Perca flavescens</u>)	24.2°C (75.6°F)	Ferguson (1958)
Muskellunge (<u>Esox masquinongy</u>)	24.0°C (75.2°F)	Jackson and Price (MS, 1949)
Burbot (<u>Lota lota lacustris</u>)	21.2°C (70.2°F)	Crossman, Ireys and Pecicock (MS, 1953)
Yellow Perch (<u>Perca flavescens</u>)	21.0°C (69.8°F)	McCracken and Starkma (MS, 1948)
Brown Trout (<u>Salmo trutta</u>)	12.4-17.6°C (54.3-63.7°F)	Tait (MS, 1958)
Brook Trout (<u>Salvelinus fontinalis</u>)	14.0-16.0°C (57.2-60.8°F)	Graham (1948)
Rainbow Trout (<u>Salmo gairdnerii</u>)	13.6°C (56.5°F)	Garside and Tait (MS, 1958)
Lake Whitefish (<u>Coregonus clupeaformis</u>)	12.7°C (54.9°F)	Tompkins and Fraser (MS, 1950)
Lake Trout (<u>Salvelinus namaycush</u>)	12.0°C (53.6°F)	McCauley and Tait (MS, 1956)

TABLE 4

FIELD OBSERVATIONS ON VARIOUS SPECIES OF FISH
AND ASSOCIATED TEMPERATURES

Some temperatures are estimates derived from Ferguson's tabled or figured data. August distributions and temperatures were used wherever possible. These figures represent the temperature of the water strata in which the fish seemed to concentrate. It is judged these represent preferred natural temperature.

(This Table Taken in Part from Ferguson, 1958)

Species	Temperature	Water	Location	Author
Bass, Largemouth (<u>Micropterus salmoides</u>)	26.6-27.7°C (80.0-81.9°F)	Norris Reservoir	Tennessee	Dendy 1948
Bass, Spotted (<u>Micropterus punctulatus</u>)	23.5-24.4°C (74.1-75.9°F)	Norris Reservoir	Tennessee	Dendy
Walleye (<u>Stizostedion v. vitreum</u>)	20.6°C (69.1°F)	Trout Lake	Wisconsin	Hile and Juday, 1941
Walleye (<u>Stizostedion v. vitreum</u>)	22.7-23.2°C (72.9-73.8°F)	Norris Reservoir	Tennessee	Dendy 1948
Gizzard Shad (<u>Dorosoma cepedianum</u>)	22.5-23.0°C (72.5-73.4°F)	Norris Reservoir	Tennessee	Dendy
Freshwater Drum (<u>Aplodinotus grunniens</u>)	21.6-22.2°C (70.9-72.0°F)	Norris Reservoir	Tennessee	Dendy
Rock Bass (<u>Ambloplites rupestris</u>)	14.7-21.3°C (58.5-70.3°F)	Lakes	Wisconsin	Hile and Juday, 1941
Rock Bass (<u>Ambloplites rupestris</u>)	20.7°C (69.3°F)	Streams	S. Ontario	Hallam 1958
Yellow Perch (<u>Perca flavescens</u>)	21.2°C (70.2°F)	Lake Opeongo	Ontario	Present Work
Yellow Perch (<u>Perca flavescens</u>)	21.0°C (69.8°F)	Costello Lake	Ontario	Present Work
Yellow Perch (<u>Perca flavescens</u> , small)	12.2°C (54.0°F)	Muskellunge Lake	Wisconsin	Hile and Juday, 1941
Yellow Perch (<u>Perca flavescens</u> , larger)	20.2°C (68.4°F)	Muskellunge Lake	Wisconsin	Hile and Juday, 1941

TABLE 4, continued

Species	Temperature	Water	Location	Author
Yellow Perch (<u>Perca flavescens</u> , larger)	20.2°C (68.4°F)	Silver Lake	Wisconsin	Hile and Juday, 1941
" "	21.0°C (69.8°F)	Nebish Lake	Wisconsin	Hile and Juday, 1941
" "	20.8°C (69.5°F)	Trout Lake	Wisconsin	Hile and Juday, 1941
" "	19.7°C (67.5°F)	Lake Nipissing	Ontario	Present Work
Bass, Smallmouth (<u>Micropterus dolomieu</u>)	20.3-21.3°C (68.5-70.3°F)	Nebish Lake	Wisconsin	Hile and Juday, 1941
" "	21.4°C (70.5°F)	Streams	S. Ontario	Hallam 1958
Sauger (<u>Stizostedion canadense</u>)	18.6-19.2°C (65.1-66.6°F)	Norris Reservoir	Tennessee	Dendy 1948
Brook Trout (<u>Salvelinus fontinalis</u>)	14.2-20.3°C (57.6-68.5°F)	Moosehead Lake	Maine	Cooper and Fuller
" "	15.7°C (60.3°F)	Streams	S. Ontario	Hallam 1958
" "	12.0-20.0°C (53.6-68.0°F)	Redrock Lake	Ontario	Baldwin 1948
Mottled Sculpin (<u>Cottus bairdii</u>)	16.5°C (61.7°F)	Streams	S. Ontario	Hallam 1958
Brook Trout x Lake Trout (<u>Salvelinus</u> hybrid)	13.1°C (55.6°F)	Jack L. Sproule Lake	Ontario	Martin and Baldwin, 1958
White Sucker (<u>Catostomus commersonnii</u>)	11.8-20.6°C (53.3-69.1°F)	Musk, Trout Silver	Wisconsin	Hile and Juday, 1941
White Sucker (<u>Catostomus</u> c. <u>commersonnii</u>)	14.1-18.3°C (57.4-64.9°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
Round Whitefish (<u>Prosopium cylindraceum</u>)	13.9-17.5°C (57.0-63.5°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
Alewife (<u>Pomolobus pseudoharengus</u>)	4.4- 8.8°C (39.9-47.8°F)	Cayuga Lake	New York	Galligan 1951

TABLE 4 , continued

Species	Temperature	Water	Location	Author
Lake Trout (<u>Salvelinus namaycush</u>)	10.0-15.5°C (50.0-59.0°F)	Cayuga Lake	New York	Galligan 1951
" "	14.0°C (57.2°F)	White Lake	Ontario	Kennedy 1941
" "	11.0-11.5°C (51.8-52.7°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
" "	8.0-10.0°C (46.4-50.0°F)	Louisa Redrock	Ontario	Martin 1952
American Smelt (<u>Osmerus mordax</u>)	12.8°C (55.0°F)	Lake Champlain	New York	Greene 1930
" "	6.6- 8.3°C (43.9-47.9°F)	Cayuga Lake	New York	Galligan 1951
Lake Whitefish (<u>Coregonus clupeaformis</u>)	11.4-11.9°C (52.5-53.5°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
Longnose Sucker (<u>Catostomus catostomus</u>)	11.0-11.6°C (51.8-53.0°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
Burbot (<u>Lota lota maculosa</u>)	10.8-11.4°C (51.4-52.5°F)	Moosehead Lake	Maine	Cooper and Fuller, 1945
Coregonys (<u>Leucichthys artedi</u>)	8.0-10.0°C (46.4-50.0°F)	Lake Nipissing	Ontario	Fry 1937
" " (Cisco or Lake Herring)	5.5- 7.2°C (41.9-44.8°F)	Cayuga Lake	New York	Galligan 1951




FIGURE 4

FIELD OBSERVATIONS OF FISH AND ASSOCIATED TEMPERATURES DURING MIDSUMMER (AUGUST MOSTLY). THE DEPTH OF EACH RECTANGLE CORRESPONDS TO THE TEMPERATURE RANGE. POINTERS ON Laterally rounded figures represent a derived average. VERTICAL RELATIONS ONLY ARE IMPORTANT (FERGUSON, 1958),




FIGURE 5. FIELD OBSERVATIONS OF FISH AND ASSOCIATED TEMPERATURES DURING MIDSUMMER (AUGUST MOSTLY). THE DEPTH OF EACH RECTANGLE CORRESPONDS TO THE TEMPERATURE RANGE. POINTERS ON Laterally rounded figures represent a derived average. VERTICAL RELATIONS ONLY ARE IMPORTANT (FERGUSON, 1958).

may not kill trout they produce environmental conditions favorable for the production of coarse fish.

One unusual set of data collected on preferred temperatures of rainbow trout by Garside and Tait (1958) showed that the preferred temperatures were inversely related to the acclimation temperature. They state, "Fish cannot lose heat because they must pass considerable quantities of water through their respiratory system in order to compensate for the low quantities of dissolved oxygen contained in waters possessing a higher temperature. With the animal passing higher quantities of heated water across their gills the body temperature of the animal must rise."

Effect of Temperature on Toxicity

Effects of artificially induced temperature changes can result in fish mortalities; as temperature increases the toxicity of certain materials increases.

The Prevention Subcommittee of the Central Water Committee, Ministry of Health, England, (Anon., 1949) states, "...increase in temperature also increases the lethal effect of toxic substances to fish." For example a rise in temperature from 8°C (46°F) to 18°C (64°F) approximately doubled the toxicity of a low concentration of potassium cyanide.

The toxicity of chloride concentrations has been shown to be dependent on temperature. The temperature has a significant effect on the time of both initial and final mortalities, the rate of mortality, and the duration

of the mortality for rainbow trout. It has been postulated that the metabolic rate of the fish, which affects the rate to which fluoride is toxic to rainbow trout, is affected by the increased temperature of the fish, Angelovic (1961).

Benefits

Trembley (1960) concluded that most fish species are attracted to and invade heated water areas from late September until early June. Attraction to heated water has been observed in England, and has been reported frequently in America. This adds to the recreational value of localized areas, because angling can be continued throughout the winter when there may be little or no fishing in other areas. Trembley found that one of the disadvantages to providing winter fishing is that fish leave the heated-water zone in the hot summer months.

Another benefit of artificially induced temperature changes is the production of trout and other cold water fish in the reach downstream from reservoirs. Low level penstock discharges from stratified reservoirs often lowers the temperature in the receiving stream to 12.8°C (55°F) and it may not exceed 20°C (68°F) even in summer (Mackenthun et al., 1964).

Summary

1. Warm water fish can survive temporarily in waters heated artificially to 33.9°C (93°F); even at 30°C (86°F) coarse fish populations, such as roach, perch, gudgeon, tench and carp, are reduced. In cold weather, stream temperature should be substantially below 33.9°C (93°F) to prevent mortalities when fish move through excessive stream gradients.
2. Streams supporting cold water non-anadromous fish populations should not receive heated effluents that will raise receiving stream temperatures above 14.5°C (58°F). In cold weather, stream temperature should be below 14.5°C (58°F) to prevent mortalities.
3. Sudden changes in temperature can be more harmful to some species of fish than continued exposure to a higher temperature.
4. Fish can adapt to higher temperatures faster than to lower temperatures.
5. The maximum temperature for a given species of fish varies with the fish's rate of heating, size, and physiological condition.
6. Fish may starve at elevated temperatures because of their inability to capture food.
7. Fish seek out a preferred temperature at which they can best survive which is several degrees below their lethal temperature.
8. The toxic effects to fish of certain materials increase with temperature.
9. Certain benefits, including open water winter fishing in otherwise ice covered areas and a cold water fisheries downstream from reservoirs, can be derived from artificially induced temperature changes. The

benefits of fish being attracted to heated water in the winter months may be negligible compared to fish mortalities that may result when the fish return to the cooler water; lethal temperatures may result from heated discharges in the summer months.

REFERENCES CITED

- Agersborg, H. P. K., 1930. The influence of temperature on fish. *Ecology*, 11(1): 136-144.
- Alabaster, J. S., 1962. The effect of heated effluents on fish. *Internat. Conf. on Water Poll. Resrch.*, London, 1962. *Air and Water Poll.* 7(6/7): 541-563, 1963. *Bio. Abstr.*, 45(13), Abstr. No. 54305, 1964.
- Angelovic, J. W., W. F. Sigler, and J. M. Neuhold. Temperature and fluorosis in rainbow trout. *Jour. Water Poll. Control Fed.*, 33: 371-381, Apr. 1961.
- Anon., 1949. Discharge of heated liquids into streams. Rep. of Rivers Poll. Prevention Sub-Committee of the Central Water Committee, Minist. of Health, England, pp. 69-76.
- Anon, 1962. Heated discharges . . . their effect on streams. Rep. by the Advisory Committee for the control of stream temperatures to the Pa. Sanitary Water Board. Pa. Dept. Health, Harrisburg, Publ. No. 3, 108 pp.
- Brett, J. R., 1960. Thermal requirements of fish - three decades of study, 1940-1970. *Bio. Problems in Water Poll.*, Trans. 1959 Seminar, Robert A. Taft Sanitary Engng. Center Tech. Rep. W60-3, Cincinnati, pp. 110-117.
- Brown, M. E., 1957. The physiology of fishes. Vol. I. Metabolism. Vol. II. Behavior. Academic Press.
- Cairns, J., Jr., 1956. Effect of heat on fish. *Indus. Wastes*, 1(5): 180-183.
- Downing, K. M., and C. J. Merckens, 1957. The influence of temperatures on the survival of several species of fish in low tensions of dissolved oxygen. *Ann. Appl. Bio.*, 45(2): 261-267.
- Ellis, M. M., 1947. Temperature and fishes. U. S. Fish and Wildl. Serv., Fish Leaflet 221.
- Falkner, N. W., and A. H. Houston, 1966. Some haematological responses to sublethal thermal shock in the goldfish, Carassius auritus L. *Jour. Fish. Resch. Bd. Can.*, 23(8): 1109-1120.

Ferguson, R. G., 1958. The preferred temperature of fish and their mid-summer distribution in temperate lakes and streams. Jour. Fish Resch. Bd. Can., 15: 607-624. Spo. Fish. Abstr., 4(1), Abstr. No. 2275, 1959.

Fry, F. E. J., 1947. Effects of the environment on animal activity. Univ. of Toronto Stud. Bio. Ser. 55: Publ. Ont. Fish. Resch. Lab., 68: 1-62.

Fry, F. E. J., J. S. Hart, and K. F. Walker, 1946. Lethal temperature relations for a sample of young speckled trout, Salvelinus fontinalis, Univ. Toronto Stud. Bio. Ser. 54, Publ. Ont. Fish Resch. Lab., 66: 1-35.

Garside, E. T., and J. S. Tait, 1958. Preferred temperature of rainbow trout (Salmo gairdneri Richardson) and its unusual relationship to acclimation temperature. Can. Jour. Zool., 36: 563. Spo. Fish. Abstr., 3(3), Abstr. No. 1676, 1958.

Heinicke, E. A., and A. H. Houston, 1965. Effect of thermal acclimation and sublethal heat shock upon ionic regulation in the goldfish Carassius auratus L., Jour. Fish Resch. Bd. Can., 22(6): 1455-1476.

Jones, J. R. E., 1964. Thermal pollution: the effect of heated effluents. Fish and River Pollution, Chap. 13: 153-168, Butterworth and Co., Ltd., Washington, D. C.

Mackenthun, K. M., W. M. Ingram, and R. Porges, 1964. Limnological aspects of recreational lakes. U. S. Dept. of Health, Educ. and Welfare, GPO 176 pp.

Prosser, C. L., 1955. Physiological variations in animals. Bio. Rev., 30(3): 229-262.

Swift, D. R., 1965. Effect of temperature on mortality and rate of development of the eggs of the Windermere Char (Salvelinus alpinus). J. Fish. Res. Bd. Canada, 22(4).

Tarzwel, C. M., 1957. Water quality criteria for aquatic life. In: Biological Problems in Water Pollution. Robert A. Taft Sanit. Engng. Center, Cincinnati, Ohio, pp. 246-272.

Threinen, C. W., 1958. Cause of mortality of a midsummer plant of rainbow trout in a southern Wisconsin lake, with notes on acclimation and lethal temperatures. Prog. Fish. Cult., 20: 27. Spo. Fish. Abstr., 3(2), Abstr. No. 1543, 1958.

Trembley, F. J., 1960. Research project on effects of condenser discharge water on aquatic life, progress report, 1956 to 1959. Inst. of Resch., Lehigh Univ., 154 pp. Water Poll. Abstr., 34(11), Abstr. No. 2157, 1961.

Wells, M. M., 1914. Resistance and reactions of fishes to temperature. Trans. Illinois Acad. Sci., 7: 48-59.

IV. MARINE, ESTUARINE AND ANADROMOUS FISHES

Introduction

Because a growing demand for electricity supplied by steam or nuclear generators has increased the need for ample water, both for cooling and thermal waste assimilation, larger areas of fresh and marine waters will receive significant temperature elevations (Mihursky and Kennedy, 1967). Future emphasis will be away from the use of inland waters, toward the utilization of estuarine and marine resources (Naylor, 1965). The protection of fishes in the estuarine and marine areas, as well as anadromous fishes that must move through areas of heated water is becoming an increasing problem.

General Temperature Effects

Researchers, studying the effects of fluctuating temperatures on fishes, have taken two approaches: one method of study is to observe the reactions of fishes in their natural habitat; the other method is to remove representative samples of fish from the natural habitat to the laboratory and observe them under selected test conditions. Both methods of study have been used equally well.

Physiology

The physiology of fishes is directly affected by temperature. Fishes are classed as poikilothermus or animals whose body temperature follows

changes in environmental temperatures rapidly and precisely. In such animals the factors favoring heat loss tend to equal the factors producing body heat and the body temperature approaches environmental temperature (Prosser, et al., 1950; Kinne, 1963). In a majority of fishes the body temperature differs from that of the surrounding water by only 0.5-1.0°C (0.9-1.8°F) (Nikolsky, 1963). Therefore, one of the fundamental requirements of fishes is that the external temperature must be best suited to internal tissues (Brett, 1956). Cells exposed to heat undergo an increase in the viscosity of protoplasm. This increase is reversible to a point beyond which heat death occurs (Gunter, 1957). Various functions of an organism, such as reproduction, locomotion and growth may have different temperature ranges and these ranges should be known in order to evaluate the effects of temperature on that organism (Kinne, 1963). For example, according to Kinne (1963) and Naylor (1965), marine and brackish water organisms may increase or decrease osmotic regulation as a function of temperature.

Temperature fluctuations act on an organism in a variety of ways:

a) metabolic rates are changed, b) reproduction is affected, c) distribution may be increased or decreased and d) tolerance limits are widened or narrowed.

Metabolism

Rates of metabolism and activity increase with increasing temperatures over most of the tolerated temperature range and then often cease suddenly near the upper lethal temperature. Such rates vary with different species,

processes and levels or ranges of temperature and may be modified by salinity and oxygen factors (Kinne, 1963). Changes in metabolic rates, because of temperature fluctuations, may be signaling factors for spawning or migration (Nikolsky, 1963).

Reproduction

The effects of temperature on reproduction in many animals are confined to narrower ranges than the majority of functions (Kinne, 1963 and Gunter, 1957). Most marine animals have restricted temperatures for breeding. Rising temperatures in the spring induce development of the gonads and actual spawning takes place when a certain temperature level is reached, which varies for different species. Some fish spawn on a drop in temperature while others respond to a rise in temperature (Gunter, 1957). Because of narrow breeding requirements the survival of a species in heated waters does not preclude the possibility that the species may be prevented from breeding and may exist in an area by continued recruitment from outside (Naylor, 1965).

Development

Temperature changes affect fish development in several ways. Meristic characters and shape may be changed as well as embryonic development. Low temperatures slow down development and in some cases many marine and brackish water animals attain a larger final size because of their slow, long continued growth rather than rapid growth (Kinne, 1963).

Distribution and Ecology

Since temperature is the most important single factor governing the occurrence and behavior of life, it not only affects the distribution of a single species, it may also modify the species composition of a community or an ecosystem (Gunter, 1957 and Kinne, 1963). Tropical and subtropical fishes are more stenothermal (tolerate a narrow range of temperatures) than those of boreal and higher latitudes and marine forms are more stenothermal than fresh water ones (Nikolsky, 1963). In his publication on temperature effects on marine organisms Naylor (1965) noted that estuarine forms were more tolerant of heated effluents than marine forms and littoral species, and concluded that some coldwater stenothermal forms may be eliminated by heated discharges and some eurythermal (tolerate a wide range of temperatures) species may be increased. He also noted that temperature effects seem to be more pronounced in enclosed areas of estuaries and bays, while heat effects in open estuaries are least striking. In tropical areas, species live close to their thermal limits and effects of heated effluents are more pronounced, while in northern (Arctic) areas species may be 13 to 16°C (23.4 to 28.8°F) below their death temperatures and may not be as severely affected.

By testing species in the laboratory Brett (1956) noted that a slow rate of decrease in environmental temperature is of greater importance for

maintaining life than a slow rate of increase. Lethal cold can be more important than lethal heat as a factor limiting the distribution of marine fish and as a hazard to some in their native habitats (Doudoroff, 1957).

Acclimation

The capacity to acclimate depends on the genetic background, environmental history and present physiological condition and age of the organism involved (Kinne, 1963). For example, the resistance of animals to cold is much more variable than resistance to heat and resistance to cold varies with size, smaller fish resisting best (Gunter, 1957).

Acclimation to different temperatures may involve changes in orientation, migration, and other behavioral aspects such as territorialism as well as biological rhythms (Kinne, 1963). In his experiments with marine fishes, Doudoroff (1957) noted that acclimatization to heat may be acquired very rapidly, the speed varying with heat. Also, brief or intermittent exposure to high temperatures can result in markedly increased resistance to heat which is not readily lost on subsequent exposure to low temperatures. However, it is the rapidity of the onset of low temperatures that probably causes death, outstripping the ability of fish to acclimate and resulting in greater mortalities that are due to cold in nature (Brett, 1960). Deaths resulting from the inability of fish to rapidly acclimate to lowering temperatures have been reported by Gunter and Hildebrand (1951) and Galloway (1951).

According to Kinne (1963) acclimation to low temperature usually tends to shift the lower thermal limits downward and acclimation to high temperatures tends to shift the upper limits upward. As a result the ability to acclimate affects the temperature ranges that a fish can tolerate.

Tolerance

The temperature tolerance of fish varies with their development, area of distribution and physiology. As noted earlier, estuarine forms are more tolerant of heated effluents than marine forms and littoral species (Naylor, 1965). However, Kinne (1963) reports that in general, the total range of temperature tolerated in the state of active life is smallest in marine forms and largest in brackish and fresh-water forms. Gradual changes are tolerated much better than sharp changes. Some species can stand a gradual change up to 30 or 35°C (86 or 95°F), but at the upper extreme, many organisms are killed by temperatures not far above those to which they are accustomed.

MARINE FISHES*

Marine fishes that inhabit the shore line areas, estuaries or bays are most often affected by temperature changes. The problem of thermal shock to pelagic (living or occurring in the open ocean) life histories is extremely critical in marine environments (Mihursky and Kennedy, 1967).

Eggs

The effects of salinity and temperature on the eggs of Pacific Cod (Gadus macrocephalus) were studied by Forrester (1964) and, Forrester and

* In this report the term marine includes estuarine species.

Alderdice (1966). In the study by Forrester, eggs were held to completion of hatching in various combinations of salinity and temperature. Maximum hatching success was in the vicinity of 19 parts per thousand (ppt) salinity and 5°C (41.0°F). Forrester and Alderdice observed that the relationship between rate of development of cod eggs and water temperature was linear at temperatures of 5-11°C (41-51.8°F). Time to 50 percent hatching ranged from 8.5 days at 11°C (51.8°F) to 17 days at 5°C (41°F). Most successful hatching occurred at the lower temperature. One of the most noted variables in the study on eggs of the American smelt (Osmerus mordax) in Maine was the large increases in mortality during extreme fluctuations in daily water temperature of as much as 7°C (12.6°F) as observed by Rothschild (1961). Striped bass eggs (Roccus saxatilis) were found to survive in constant fluctuations of water temperatures ranging from 12.8-23.9°C (55-75°F) Albrecht (1964). The tolerance of eggs of four marine fishes was studied by Hubbs (1965). California killifish (Fundulus parvipinnis), topsmelt (Atherinops affinis), California grunion (Leuresthes tenuis) and mussel blenny (Hypsoblennius sp.) were incubated at a variety of temperatures. Larvae successfully hatched at temperatures between 16.6°C and 28.5°C (61.9-83.1°F), 12.8 (-)°C and 26.8°C (55-80.1°F), 14.8° C and 26.8°C (58.6-80.1°F) and 12.0 (-) and 26.8 (+)°C (53.6(-)-80.1(+))°F respectively (Table 5).

Young

Larvae of some marine fish are pelagic in the early part of their life history, and temperature of the surrounding water determines the rate of

Table 5 - Temperature ranges reported for the hatching of eggs from various species of marine and anadromous fish.

Species	Lower Temp. (C)	Upper Temp. (C)	Remarks	Source
<u>Roccus</u>				
<u>saxatilis</u>	12.8	23.9	Survived Constant Fluctuations	Albrecht, 1964
<u>Fundulus</u>				
<u>parvipinnis</u>	16.6	28.5	Successful Hatching	Hubbs, 1965
<u>Atherinops</u>				
<u>affinis</u>	12.8	26.8	" "	" "
<u>Leuresthes</u>				
<u>tenuis</u>	14.8	26.8	" "	" "
<u>Hypsoblennius</u>				
<u>sp.</u>	12.0(-)	26.8(+)	" "	" "
<u>Petromyzon</u>				
<u>marinus</u>	15.0	25.0	Survival Range	McCauley, 1963
"	15.6	21.1	" "	Piavis, 1961
<u>Oncorhynchus</u>				
<u>nerka</u>	4.4-5.8	12.8-14.2	Threshold Temp.	Combs, 1965
<u>O. tshawytscha</u>	5.8	14.2	" "	Combs & Burrows, 1957
" "	9.4	14.4	" "	Seymour, 1956
" "	5.6	14.4	" "	Leitritz, 1962
All Salmon	5.8	12.8	" "	Anon., 1966

development from a pelagic form to an actively swimming form. The rate of development of the lemon sole (Parophrys vetulus) determines the number of young that reach the nursery grounds in Hecate Strait, British Columbia (Ketchen, 1956). Small annual differences in the temperature of sea water produce marked differences in the duration of the pelagic stage. Below average temperatures result in the larvae being carried by the currents for a longer period of time and more larvae are deposited on the nursery ground. Thus, temperature may govern the strength of a year class. A temperature of 6.2°C (43.1°F) seems to produce the best deposition of larvae. Increases in temperature increase the rate and shorten the time of development of herring (Clupea pallasii). A temperature increase from $4.4\text{--}10.7^{\circ}\text{C}$ ($40\text{--}51.1^{\circ}\text{F}$) shortened development time from 40 to 11 days (Blaxter, 1963).

In the estuarine environment, fishes are more susceptible to heat changes. However, as noted in the general discussion, fish in estuarine waters seem to tolerate a wider range of temperatures. Striped bass fingerlings (Roccus saxatilis) were able to tolerate 35°C (95°F) in laboratory tests (Talbot, 1966). According to Talbot (1966), Merriman (1941) studied the striped bass of the Atlantic coast and found the maximum temperature in the New England area to be 25°C to 27°C (77.0 to 80.6°F) with fish kills occurring at these temperatures. Juvenile striped bass have survived transfer between salt and fresh-water at temperatures in the

range of 12.8-21.1°C (55-70°F) but they are not tolerant to changes from fresh-water to saltwater at 7.2°C (45°F) (Tagatz, 1961). Striped bass acclimated at 4.4°C (40°F) and tested for eight hours with increases in temperature of 2.3°C (3.5°F) at one hour intervals, had a median lethal dose (LD₅₀) of 23.9°C (75°F), (Trembley, 1960).

The larvae of Atlantic Menhaden (Brevoortia tyrannus) were able to survive longer when acclimated at cooler temperatures, than when acclimated to warmer temperatures. Acclimation temperature was more important to larval survival at test temperatures below 5.0°C (41°F) than at 5.0°C (41°F) and above. Larvae acclimated at 7 and 10°C (44.6 and 50°F) survived over twice as long at 4.5°C (40.1°F) as those acclimated at 12.5 or 15°C (54.5 or 59°F), (Lewis, 1965). The effects of salinity on temperature tolerances were checked by Lewis (1966), who found a temperature of 6°C (42.8°F) and below with zero ppt salinity lowered larval survival time to only a few hours. At a salinity of 5-30 ppt and a temperature of 4°C (39.2°F) larval survival was good. Lower and upper limits of salinity tolerance were increased with increasing temperature.

The relation of menhaden (B. tyrannus) to estuaries was studied by Reintjes and Pacheco (1966). A water temperature of 3°C (37.4°F) may be critical to larval survival. Larval menhaden can suffer mass mortalities when water temperatures fall below 3°C (37.4°F) for several days or chill rapidly. The matter of chill seems to be very important to estuarine fishes.

Doudoroff (1942) studied the resistance and acclimation of marine fishes to temperature changes. Young greenfish (Girella nigricans) were used in his tests. Heat resistance was gained rapidly and lost slowly. Resistance to chilling was lost slowly on warming and acquired no more rapidly on cooling. The changes of resistance to heat and to cold were found to be more or less independent and distinct phenomena. Acclimation to cold is slow, to heat, fairly fast. For these reasons injury by chilling is no less important as a possible limiting factor in the distribution of marine fishes than heat injury.

More active fishes are able to avoid harmful temperatures and to exercise selection in experimental gradient (Doudoroff, 1938). In the case of pelagic marine larvae, circumstances may dictate their survival. A mortality of marine fish larvae was noted in an area off Georges Bank. Currents carried the larvae from cold 7.8°C (46°F) water into warm layers of 20°C (68°F) water and large mortalities resulted (Colton, 1959).

A study of the effect of extreme temperatures on herring larvae (C. harengus) revealed an upper lethal temperature 22.0 to 24.0°C (71.6 - 75.2°F) and a lower lethal temperature of -0.75 to -1.8°C (30.6 - 29.1°F). Larvae were 6-8 millimeters long and acclimated to temperatures between 7.5 and 15.5°C (58.2 - 72.9°F) (Blaxter, 1960). Young topsmelt (Atherinops affinis) acclimated to temperatures of 20°C (68°F) had an upper 48 hour median tolerance limit (TL_m) of 31.8°C (89.1°F) and a lower 48 hour TL_m of 10.1°C (50.1°F), (Doudoroff, 1945). Young greenfish (Girella nigricans) acclimated to

temperatures of 12-28°C (53.6-82.4°F) exhibited a lower 48 hour TL_m of 4.1 to 13°C (39.4-55.4°F) and a upper TL_m of 28.7 to 31.5°C (81.9-88.8°F), (Doudoroff, 1942). Temperature ranges reported for young fish are listed in Table 6.

Adults

Adult fish are usually able to select their preferred temperatures, unless they are trapped in shallow waters or forced to migrate through heated or chilled areas. Fish kills have been reported in areas of shallow water. Atlantic round herring (Etrumeus sadina) and chum mackerel (Scomber colias) were observed dead and dying after several days of cold weather had dropped water temperatures in Pamlico Sound, North Carolina, to 5.2°C (41.4°F), (Wells, Wells and Gray, 1961). The effects of winter water conditions were also observed by Schwartz (1964). He noted that most fish sank when killed by lowered temperatures and would probably not be observed. In the area of high temperatures, alewives (Alosa pseudoharengus) died of heat shock after being herded into water of 26.7 - 32.2°C (80-90°F). The same species showed no effects when they entered a lagoon with 22.8°C (73°F) water (Trembley, 1960). Herring, in nature, have been found in almost all temperature ranges permitted by their resistance to temperature extremes. Herring at the appropriate season have an upper lethal temperature of 19.5 to 21.2°C (67.1 to 70.1°F) depending upon size and can survive short exposure to temperatures below -1.0°C (30.2°F), (Brawn, 1960).

Table 6 - Temperature ranges reported for young marine and anadromous fish.

Species	Acclimation Temp. (C)	Lower Lethal Temp. (C)	Upper Lethal Temp. (C)	Remarks	Source
<u>O. tshawytscha</u>	23	7.4		Approx. 4 days	Brett, 1952
	20		25.1	Approx. 7 days	"
<u>O. kisutch</u>	23	6.4		Approx. 4 days	"
	20		25.0	" 7 "	"
<u>O. nerka</u>	23	6.7		Approx. 4 days	"
	20		24.4	" 7 days	"
<u>O. gorbuscha</u>			23.9	Approx. 7 days	
<u>O. keta</u>	23	7.3		Approx. 4 days	"
	20		23.8	Approx. 7 days	"
<u>Clupea</u>					
<u>harengus</u>	7.5-15.5	-0.75to-1.8	22.0-24.0		Blaxter, 1960
<u>Atherinops</u>					
<u>affinis</u>	20	10.1	31.8	48 hr TL _m	Doudoroff, 1942b
<u>Girella</u>					
<u>nigricans</u>	12-28	4.1-13	28.7-31.5	48 hr TL _m	Doudoroff, 1942a

Adult white perch (Roccus americana) acclimated to 4.4°C (40°F) and tested for 8 hours with increases of 2.0°C (3.5°F) every hour exhibited a median lethal dose (LD_{50}) of 27.8°C (82.0°F), (Trembley, 1960). In a flowing water test with heat increases of 1.1°C (2°F) per hour saltwater killifish (Fundulus heteroclitus) acclimated at 7.2°C (45°F) had a LD_{50} of 37°C (99°F) (Trembley, 1961). Adult California killifish (F. parvipinnis) were tested by Doudoroff (1945). At acclimation temperatures of $14\text{--}28^{\circ}\text{C}$ ($57.2\text{--}82.4^{\circ}\text{F}$) the upper TL_m 's, for 48 hour tests, were 32.3 to 36.5°C ($90.1\text{--}97.7^{\circ}\text{F}$). The lower 48 hour TL_m was 30°C (86°F) for fish acclimated at 20°C (68°F). Striped bass occur in wide ranges of temperatures in the estuary (Talbot, 1966). They will spawn between 14.4°C (58°F) and 21.1°C (70°F). Ranges of temperature tolerated are $6.0\text{--}7.5^{\circ}\text{C}$ ($42.8\text{--}45.5^{\circ}\text{F}$) to $25\text{--}27^{\circ}\text{C}$ ($77\text{--}80^{\circ}\text{F}$). Temperature tolerances of three marine fishes have been determined by Hoff and Westman (1966). The common silverside (Menidia menidia) acclimated at temperatures ranging from $7\text{--}28^{\circ}\text{C}$ ($44.6\text{--}82.4^{\circ}\text{F}$) had an upper 48 hour TL_m range of 22.5 to 32.5°C ($72.3\text{--}90.3^{\circ}\text{F}$) and a lower range of 1.5 to 8.7°C ($34.8\text{--}47.8^{\circ}\text{F}$). Winter flounder (Pseudo pleuronectes) acclimated at temperatures of $21\text{--}28^{\circ}\text{C}$ ($69.8\text{--}82.4^{\circ}\text{F}$) had a lower 48 hour TL_m range of $1.0\text{--}5.4^{\circ}\text{C}$ ($33.8\text{--}41.6^{\circ}\text{F}$). Flounder acclimated at temperatures from $7\text{--}28^{\circ}\text{C}$ ($44.6\text{--}82.4^{\circ}\text{F}$) had an upper range of $22\text{--}29^{\circ}\text{C}$ ($71.6\text{--}84.2^{\circ}\text{F}$). Northern swellfish (Spheroides maculatus) were acclimated at temperatures of $14\text{--}28^{\circ}\text{C}$ ($57.2\text{--}82.4^{\circ}\text{F}$) and had a lower 48 hour TL_m of $8.4\text{--}13^{\circ}\text{C}$ ($47.1\text{--}55.4^{\circ}\text{F}$). Fish acclimated at temperatures of $10\text{--}28^{\circ}\text{C}$ ($50\text{--}82.4^{\circ}\text{F}$) had TL_m 's of $28.2\text{--}33.0^{\circ}\text{C}$ ($82.9\text{--}90.4^{\circ}\text{F}$). Temperature ranges for adults are listed in Table 7.

Table 7 - Temperature ranges reported for adult
marine and anadromous fish.

Species	Acclimation Temp. (C)	Lower Lethal Temp. (C)	Upper Lethal Temp. (C)	Remarks	Source
<u>Clupea harengus</u>		- 1.0	19.5-21.2		Brawn, 1960
<u>Fundulus</u>					
<u>parvipinnis</u>	14-28		32.3-36.5	48 hr TL _m	Doudoroff, 1942b
"	20	30		" " "	" "
<u>Roccus saxatilis</u>		6.0-7.5	25-27	Tolerated in Estuary	Talbot, 1966
<u>Menidia menidia</u>	7-28	1.5-8.7	22.5-32.5	48 hr TL _m	Hoff & Westman, 1966
<u>Pseudo pleuronectes</u>	21-28	1.0-5.4		48 hr TL _m	"
	7-28		22-29	" " "	"
<u>Spheroides</u>					
<u>maculatus</u>	14-28	8.4-13		" " "	"
	10-28		28.2-33.0	" " "	"
Adult Salmon		0.0	26.7	Survival Temp.	Anon., 1966

In a statement about the observed presence or absence of fish in heated areas, Trembley (1960) reported that fish may be eliminated from heated zones during warm months, but may congregate in heated areas during winter months.

ANADROMOUS FISHES

Anadromous fish are unique in their life histories. Eggs incubate in fresh water and the resulting young spend a period of a few months to several years in fresh water, then migrate to saltwater, where they grow into adults. As adults the fish mature in salt water and return to fresh water to spawn. During their life cycle, anadromous fish are subjected to various stresses such as salinity (osmotic) change, physical change, predators, and temperature (Brett, 1957).

Eggs

Investigations have shown that thermal requirements in the very early stages are more exacting than in the adult (Brett, 1956). Eggs of the sea lamprey (Petromyzon marinus) require the most exacting thermal levels (McCauley, 1963). The range of constant temperatures necessary for successful hatching is narrow, being 15-25°C (59-77°F). The range could be extended to 12-26°C (53.6-78.8°F) if the eggs were able to develop to the head stage before they were subjected to increased temperatures. Similar results were noted by Piavis (1961) who reared sea lamprey eggs at low constant temperatures and was unable to grow viable burrowing larvae at any temperature below 15.6°C (60°F) or above 21.1°C (70°F). Constant temperatures were used in the

incubation of chinook and sockeye salmon eggs (Oncorhynchus tshawytscha and O. nerka) (Combs, 1965). Chinook salmon eggs which had developed to the 128 cell stage could tolerate 1.7°C (35°F) water for the remainder of the incubation period. Sockeye salmon eggs were less resistant to high temperatures and more resistant to cold temperatures. Their lower threshold temperatures for normal development were $4.4\text{--}5.8^{\circ}\text{C}$ ($40\text{--}42.5^{\circ}\text{F}$) and upper threshold temperatures were $12.8\text{--}14.2^{\circ}\text{C}$ ($55\text{--}57.5^{\circ}\text{F}$). A lower threshold for chinook eggs was established at 5.8°C (42.5°F) and an upper threshold at 14.2°C (57.5°F), (Combs and Burrows, 1957). Mortalities occurred when eggs were incubated above or below these temperatures.

Hayes, in 1949, subjected salmonid eggs to extreme temperatures and noted that certain tissues will exhibit cell multiplication without differentiation. Salmon embryos were incubated by Hayes, Pelluet and Gorhan (1953) in temperatures within the limits for survival. Hatching of the embryos tended to appear precociously at low temperatures.

According to Johnson and Brice (1953) reservoir water could be used for incubation when the daily mean temperatures were below 12.2°C (54°F). Chinook eggs incubated over 15.6°C (60°F) suffered excessive mortality. Results from laboratory tests conducted by Olson and Foster (1957) and Nakatani and Foster (1966) were slightly higher. They reported that chinook eggs, especially in cold water, could begin incubation at temperatures as high as 16.1°C (61°F), without significant loss. Seymour (1956) reported that young chinook eggs

should be reared at temperatures ranging from 9.4°C to 14.4°C (49 to 58°F) for best results. Abnormal fry and the hatching period were increased by higher and lower temperatures. A slightly wider range of temperatures 5.6-14.4°C (42-58°F) was suggested by Leitritz (1962). As a general range for all Pacific salmon eggs LT_{50} 's were reported at 2.5°C (36.5°F) and 16.0°C (60.8°F) but less than normal survival was noted below 5.8°C (42.5°F) and above 12.8°C (55°F), (Anon, 1966).

In the natural environment, McNeil (1966) studied the effect of low temperatures in the spawning beds of pink (O. gorbuscha) and chum (O. keta) salmon and determined that freezing was important only when the maximum day-time temperatures remained below 0°C (32°F) for at least two days. Eggs and larvae of pinks and chum are able to survive at low temperatures and high salinities (Rockwell, 1956).

In a summary of the significance of temperatures on salmon egg incubation (Anon, 1966) the following points were emphasized; a) the effects of temperature vary with many things including species and race, b) mortality attributable to temperature is also a function of duration of exposure, c) temperature during the initial incubation period is critical and, d) if the initial incubation temperatures are below 5.6°C (42°F) or above 12.8°C (55°F) less than normal survival can be expected. Temperature ranges reported for the hatching of eggs are tabulated in Table 5.

Young

Young of anadromous fish, especially salmon, spend much of their life in fresh-water and most research on them pertains to this environment. Determination of tolerance limits for different species of fish may be very difficult and time consuming. Brett (1960) has reported the ultimate upper lethal levels can differ between species by as much as 17°C (31°F).

Five species of Pacific Coast salmon were tested by Brett (1952) for their temperature tolerances. The five species of salmon tested were spring or chinook (Oncorhynchus tshawytscha), silver (O. kisutch), pink (O. gorbuscha), sockeye (O. nerka) and chum (O. keta). Fish used in the tests were less than one year of age with an average length of 405 centimeters and an average weight of 1 gram. The maximum acclimation temperature was 24°C (75.2°F). Springs were reported to be very active and good feeders at 24°C (75.2°F) but growth was poor. Pinks, sockeyes, chum and cohos were all intolerant to 24°C (75.2°F) water. Of the species tested, springs and cohos were most tolerant to prolonged high temperatures, sockeyes intermediate, and pink and chum least tolerant. The upper lethal temperatures were as follows: spring - 25.1°C (77.2°F); coho - 25.0°C (77.0°F); sockeye - 24.4°C (76.0°F); pink - 23.9°C (75.0°F); and chum - 23.8°C (74.8°F). Acclimation temperatures for all species were $10\text{-}20^{\circ}\text{C}$ ($50\text{-}68^{\circ}\text{F}$). The lower lethal temperatures for the highest acclimation of 23°C (73.4°F) were spring, 7.4°C (45.4°F); coho, 6.4°C (43.5°F); sockeye, 6.7°C (44.0°F); and chum, 7.3°C (45.1°F). For all species the region of greatest preference was $12\text{-}14^{\circ}\text{C}$

(53.6-57.2°F), (Table 6). In a report on the Columbia River salmon (Anon., 1956) lethal tolerances for 50 percent of the juveniles tested were listed as 0.0°C (32.0°F) and 25.1°C (77.2°F). However, poor growth was reported for temperatures below 4.4°C (40°F) and above 18.3°C (65°F). In contrast with the above results Kerr (1953) tested young chinook salmon and reported a maximum temperature of 26.7°C (80°F) tolerated by them. He also reported them able to tolerate a rise in temperature of 9°C (16°F) in the cooling water from a steam generated electric plant.

Burrows (1963) suggests that a range in temperature for maximum productivity in fingerlings should be between 10-15.6°C (50-60°F). His research indicated that to attain maximum productivity the water temperature must not only remain within the tolerance level of the fingerling but in species with more than a minimum stay in fresh-water, the temperature must reach that necessary for optimum growth level as well.

Tests designed to reveal the effects of temperature on the physiology of young salmon showed internal temperatures of smolts reached equilibrium with the external environment in 3-5 minutes (Harvey, 1964). Effects of temperature on fin ray and vertebrae counts were checked by Seymour (1959) who noted that the average number of vertebrae per lot of fish was less for lots reared at temperatures in the middle portion of the 3.9-16.7°C (39°F-62°F) range than for lots reared at either extreme of the range.

Several researchers have checked the effects of temperature on swimming speed and metabolism. Optimum cruising speeds occurred at 15°C (59°F) for young sockeye and 20°C (68°F) for young coho (Brett, Hollands and Alderdice, 1958). When young sockeye salmon swimming at a speed of 1 foot per second were subjected to a temperature change of 10-15°C (18°F to 27°F) their metabolic rate increased by more than 50 percent (Anon., 1962).

Migration of young fish from fresh-water to saltwater may subject them to wide ranges of temperature. Effects of temperature on pink salmon (O. gorbuscha) were studied by Sheridan (1960 and 1961). He reported an interaction between air temperature and snow that may cause fry to migrate to sea at unfavorable times. There is presumably one "best" time for fry to enter saltwater and the normal time of seaward migration may be best for food supplies and saltwater temperatures or other unknown factors. In addition to the temperature and stream flow in the river, the temperature and salinity of the marine environment during early life are also very important (Vernon, 1958).

During their stay in fresh-water young fish are subjected to diseases that may or may not be influenced by temperature. Both the literature and research show the effects of some diseases are increased by temperature increases. There is one disease, however, that may be reduced by increased temperature (Ordal and Pacha, 1963). The myxobacterium Cytophaga psychrophila is a disease of salmon fingerlings in low temperature water. Losses due to

this disease can be reduced by increasing temperatures above 6.1°C (43°F) in the spring. Water temperature remaining relatively stable either above or below the optimum range for extended periods is conducive to disease development which may result in reduction of fingerlings produced (Burrows, 1963).

Adults

Adult anadromous fishes spend most of their life cycle in the marine area and enter fresh-water only to migrate up a stream to spawn. During their migration salmon do not take in food after reaching the estuary and heading upstream. High water temperatures increase their metabolic rate and may result in fuel depletion before the fish can spawn (Anon., 1962). Temperature was listed as one of the factors affecting timing of spawning runs of pink salmon (O. gorbuscha) studied by Sheridan (1960). For pinks, a stream temperature near 10°C (50°F) seemed to be best. Adult salmon have been reported to survive 0.0°C (32.0°F) to 26.7°C (80°F) but spawning effectiveness may be reduced below 7.2°C (45°F) and above 15.6°C (60°F) (Anon., 1966), (Table 7).

The migration of fish has been hampered by unfavorable temperatures. Brett (1957) noticed the curtailment of the migration of sockeye salmon through lakes in the spring and Major and Mighell (1966) concluded that rising or stable temperatures above 21.1°C (70°F) tended to block the entry of migrating fish from the Columbia River to the Okanogan River. A study, by Massmann (1957), of the relationship of water temperature and shad catches

in the York River for a three year period showed that greatest catches-per-unit effort were made at water temperatures of 7.2°C (45°F) to 15°C (59°F). Below a water temperature of 4.4°C (40°F) the fish stopped migrating and no catches were made. The death of migrating Atlantic salmon (Salmo salar) in Nova Scotia, due to low water and high temperatures was recorded by Huntsman (1942). Fresh run grilse died at about 29.5°C (85.1°F) and acclimated grilse at about 30.5°C (86.8°F).

As noted earlier with juvenile salmon, temperature increases usually result in an increase in disease which lowers the surviving numbers of spawning fish.

Summary

Temperature fluctuations affect the metabolism, reproduction, distribution, ecology and tolerance of fishes. The effect of a fluctuation depends on the species of fish, the stage in the life history of the fish, the rate of decrease or increase in temperature and the amount of thermal fluctuation. In the marine environment temperature changes are most important in enclosed areas such as the estuaries and bays as opposed to open areas. Tolerance to temperature fluctuations is least in marine forms and greatest in estuarine and fresh-water forms.

Pelagic forms are most susceptible to temperature fluctuations since they are dependent upon water currents for much of their movement. Adult fish are usually able to select their preferred temperature gradient unless trapped in shallow or enclosed areas or forced to migrate through heated or chilled areas. Most fish have restricted ranges of temperature within which they can reproduce successfully. Larval development also requires narrow ranges of temperature. For these reasons a fish population may exist in a heated area only by continued recruitment from the outside. In such areas fish may be absent during warm summer months and present in cold winter months. In some areas populations of widely tolerant species may replace stenothermal species.

Increasing temperatures may block the migrations of anadromous fish and increase the effects of diseases on those fish. However, there are reports

of temperature increases reducing losses of young salmon from disease and increasing survival of eggs.

Cold is as important to fish populations as heat because of the inability of fish to acclimate quickly to rapid decreases in temperature. Thus, in some areas fish populations may be limited by decreases as well as increases in temperature.

REFERENCES CITED

- Albrecht, A. B., 1964. Some observations on factors associated with survival of striped bass eggs and larvae. *California Fish & Game Jour.*, 50: 2, 100-113.
- Anon., 1962. Annual report of the Fisheries Research Board of Canada, 1961-62, for the Fiscal year ended March 31, 1962, 206 pp., Water Poll. Abstr. (Brit.), 36, 889, 1963.
- Anon., 1966. Columbia River water temperature conditions and research requirements. Water Supply and Water Poll. Control Subcommittee, Task Force on Water Temperature Evaluation, Columbia Basin Interagency Committee, v-i to v-3.
- Blaxter, J. H. S., 1960. The effect of extremes of temperature on herring larvae. *Jour. Mar. Bio. Assoc. Unit. King.*, 39, 605-608.
- Blaxter, J. H. S., 1963. The behavior and physiology of herring and other clupeids. In: *Advances in Marine Biology*, F. S. Russel (ed.), Academic Press, New York, 261-393.
- Brawn, V. M., 1960. Temperature tolerance of unacclimated herring (Clupea harengus L.), *Jour. Fish. Res. Bd. Canada*, 17(5): 721-723.
- Brett, J. R., 1952. Temperature tolerance in young Pacific salmon, genus Oncorhynchus. *Jar. Fish. Resrch. Bd. Can.*, 9(6): 265-323.
- Brett, J. R., 1956. Some principles in the thermal requirements of fishes. *Quarterly Rev. of Bio.*, 31(2): 75-87.
- Brett, J. R., 1957. Salmon research and hydroelectric power development. *Jour. Fish. Resrch. Bd. Can.*, Bull. No. 114, 26 pp.
- Brett, J. R., 1960. Thermal requirements of fish - three decades of study, 1940-1970. *Bio. Problems in Water Poll.*, Trans. 1959 Seminar, Robert A. Taft Sanitary Engng. Center Tech. Rep. W60-3, Cincinnati, pp. 110-117.
- Brett, J. R., Hollands, M., and Alderdice, D. F., 1958. The effect of temperature on the cruising speed of young sockeye and coho salmon. *Jour. Fish. Resrch. Bd. Can.*, 15(4): 587-605.

- Burrows, R. E., 1963. Water Temperature requirements for maximum productivity of salmon. In: Water temperature influences, effects and control. Proc. of the Twelfth Pacific Northwest Symposium on Water Poll. Resrch., Pacific Northwest Water Laboratory, U. S. Public Health Service, Corvallis, Oregon, pp. 29-38.
- Colton, J. B., 1959. A field observation of mortality of marine fish larvae due to warming. Limnol. and Oceanogr., 4(2): 219-222.
- Combs, B. D., 1965. Effect of temperature on development of salmon eggs. Prog. Fish. Cult., 27(3): 134-137.
- Combs, B. D. and R. E. Burrows, 1957. Threshold temperatures for the normal development of chinook salmon eggs. Prog. Fish. Cult., 19(1): 3-6.
- Doudoroff, P., 1938. Reactions of marine fishes to temperature gradients. Biol. Bull., 75: 494-509.
- Doudoroff, P., 1942. The resistance and acclimation of marine fishes to temperature changes. I. Experiments with Girella nigricans (Ayres). Bio. Bull., 83: 219-244.
- Doudoroff, P., 1945. The resistance and acclimatization of marine fishes to temperature changes. II. Experiments with Fundulus and Atherinops. Bio. Bull., 88(2): 194-206.
- Doudoroff, P., 1957. Water quality requirements of fishes and effects of toxic substances. In: The Physiology of Fishes (M. E. Brown, editor), 2: 403-430, Academic Press, Inc., New York, 503 p.
- Forrester, C. R., 1964. Laboratory observations of embryonic development and larvae of the pacific cod (Gadus macrocephalus Tilesius). Jour. Fish. Resrch. Bd. Can., 21(1): 9-16.
- Forrester, C. R., and D.F. Alderdice, 1966. Effects of salinity and temperature on embryonic development of the pacific cod (Gadus macrocephalus). Jour. Fish. Resrch. Bd. Can., 23(3): 319-340.
- Galloway, J. C., 1951. Lethal effects of the cold winter of 1939/40 on marine fishes at Key West, Florida. Copeia, 2: 118-119.
- Gunter, G., 1957. Temperature. Chapter 8. In: Treatise on Marine Ecology and Palaeoecology, I. (Ed. by J. W. Hedgpeth). Geol. Soc. Amer. Mem. 67, 159-184.

- Gunter, G., and H. H. Hildebrand, 1951. Destruction of fishes and other organisms on the south Texas coast by the cold wave of January 28-February 3, 1951. *Ecol.*, 32(4): 731-735.
- Harvey, H. H., 1964. Dissolved nitrogen as a tracer of fish movements. *Verh. int. Ver. Limnol.*, 1962, 15, 947-951. *Water Poll. Abstr. (Brit.)*, 38, 878, 1965.
- Hayes, F. R., 1949. The growth, general chemistry, and temperature relations of salmonid eggs. *Quart. Rev. Bio.*, 24(4): 281-308.
- Hayes, F. R., D. Pelluett, and E. Gorham, 1953. Some effects of temperature on the embryonic development of the salmon (Salmo salar). *Can. Jour. Zoology*, 31(1): 42-51. *Water Poll. Abstr. (Brit.)*, 26, 7, 1953.
- Hoff, J. G., and J. R. Westman, 1966. The temperature tolerances of three species of marine fishes. *Jour. Marine Res.*, 24(2): 131-140.
- Hubbs, C., 1965. Developmental temperature tolerance and rates of four southern California fishes, Fundulus parvipinnis, Atherinops affinis, Leuresthes tenuis, and Hypsoblennius sp. *Calif. Fish and Game*, 51(2): 113-122.
- Huntsman, A. G., 1942. Death of salmon and trout with high temperature. *Jour. Fish. Resch. Bd. Can.*, 5(5): 485-501.
- Johnson, H. E., and R. F. Brice, 1953. Effects of transportation of green eggs, and of water temperature during incubation, on the mortality of chinook salmon. U. S. Fish and Wildlife Service, *Prog. Fish. Cult.*, 15(3): 104-108.
- Kerr, J. E., 1953. The fish rescue project at the Pacific Gas and Electric Company's Contra Costa steam plant. *Proc. Am. Soc. Civ. Engrs.*, 79(264): 10 pp. *Water Poll. Abstr. (Brit.)*, 27, 8, 1954.
- Ketchen, K. S., 1956. Factors influencing the survival of the lemon sole (Parophrys netulus, Girard) in Hecate Strait, British Columbia. *Jour. Fish. Resch. Bd. Can.*, 13(5): 647-694.
- Kinne, O., 1963. The effects of temperature and salinity on marine and brackish water animals. I. Temperature. *Oceano. Mar. Bio. Ann. Rev.*, 1: 301-340.
- Leitritz, E., 1962. Trout and salmon culture. State of California, Department of Fish and Game, *Fish Bull. No. 107*, 169 pp.

- Lewis, R. M., 1965. The effect of minimum temperature on the survival of larval Atlantic menhaden, Brevoortia tyrannus. Trans. Am. Fish. Soc., 94(4): 409-412.
- Lewis, R. M., 1966. Effects of salinity and temperature on survival and development of larval Atlantic menhaden, Brevoortia tyrannus. Trans. Am. Fish. Soc., 95(4): 423-426.
- Major, R. L., and J. L. Mighell, 1966. Influence of Rockey Reach Dam and the temperature of the Okanogan River on the upstream migration of sockeye salmon. U. S. Fish & Wildlife Service, Fish. Bull., 66(1): 131-147.
- Massmann, W. H., and A. L. Pacheco, 1957. Shad catches and water temperatures in Virginia. Jour. Wildlife Mgmt., 21(3): 351-352.
- McCauley, R. W., 1963. Lethal temperatures of the developmental stages of the sea lamprey, Petromyzon marinus L. Jour. Fish. Resch. Bd. Can., 20, 483.
- McNeil, W. J., 1966. Effect of the spawning bed environment on reproduction of sink & chum salmon. U. S. Fish & Wildlife Service, Fish. Bull., 65(2): 495-523.
- Merriman, D., 1941. Studies on the striped bass (Roccus saxatilis) of the Atlantic coast. U. S. Fish & Wildlife Service, Fish. Bull. 50(35): 77 pp.
- Mihursky, J. A., and V. S. Kennedy, 1967. Water temperature criteria to protect aquatic life. Symposium on water quality criteria to protect aquatic life. Am. Fish. Soc., Special Publ. No. 4, pp. 20-32.
- Nakatani, R. E., and R. F. Foster, 1966. Hanford temperature effects on Columbia River fishes. Mimeo., 15 p.
- Naylor, E., 1965. Effects of heated effluents on marine and estuarine organisms. In: Advances in Marine Biology, Sir Frederick S. Russell, Editor, Academic Press, 3: 63-103.
- Nikolsky, G. V., 1963. The ecology of fishes. Academic Press, New York, 352 p.
- Olson, P. A., and R. F. Foster, 1957. Temperature tolerance of eggs and young of Columbia River chinook salmon. Trans. Am. Fish. Soc., Eighty-fifth Annual Meeting, 1955, p. 203-207, Spo. Fish. Abstr., 3, 1387, 1958.

Ordal, E. J., and R. E. Pacha, 1963. The effects of temperature on disease in fish. In: *Water Temperature Influences, Effects and Control*. Proc. of the Twelfth Pacific Northwest Symposium on Water Poll. Resch., Pacific Northwest Water Laboratory, U. S. Public Health Service, Corvallis, Oregon, pp. 39-56.

Piavis, G. W., 1961. Embryological stages in the sea lamprey and effects of temperature on development. U. S. Fish & Wildlife Service, Bu. of Commercial Fisheries, Fish. Bull. 182, 61: 111-143, Sport Fish. Abstr.. 6: 3, 1961.

Prosser, C. L., F. A. Brown, D. W. Bishop, T. L. John, and V. J. Wulff, 1950. *Comparative animal physiology*. W. B. Saunders Co., Phila., Pa., 888 pp.

Reintjes, J. W., and A. L. Pacheco, 1966. The relation of menhaden to estuaries. In: *A Symposium on Estuarine Fisheries*, Am. Fish. Soc., Special Publ. No. 3, pp. 50-58.

Rockwell, J., Jr., 1956. Some effects of sea water and temperature on the embryos of the Pacific salmon, Oncorhynchus gorbuscha (Walbaum) and Oncorhynchus keta (Walbaum). Dissertation Abstr., 16(5): 880; Spo. Fish. Abstr., 2, 1, 1956.

Rothschild, B. J., 1961. Production and survival of eggs of the American smelt, Osmerus mordax (Mitchill), in Maine. Trans. Am. Fish. Soc., 90(1): 43-48.

Schwartz, F. J., 1964. Effects of winter water conditions on fifteen species of captive marine fishes. Am. Midl. Nat., 71(2): 434-444.

Seymour, A. H., 1956. Effects of temperature upon young chinook salmon. Ph.D. Thesis, Univ. Wash., Seattle, 127 pp. Diss. Abstr. 16(11): 2249. Spo. Fish. Abstr., 2: 953, 1957.

Seymour, A., 1959. Effects of temperature upon the formation of vertebrate and fin rays in young chinook salmon. Trans. Am. Fish. Soc., 88: 58-69.

Sheridan, W. L., 1960. Relation of stream temperatures to timing of pink salmon escapements in Southeast Alaska. Symposium on Pink Salmon, H. R. MacMillan Lectures in Fisheries. Univ. of British Columbia, Vancouver, Canada, pp. 87-102

Sheridan, W. L., 1961. Temperature relationships in a pink salmon stream in Alaska. Ecology, 42: 91-98.

- Tagatz, M. E., 1961. Tolerance of striped bass and American shad to changes in temperature and salinity. U. S. Fish & Wildlife Service, Spec. Sci. Rept.-Fish., 388, 8 p.
- Talbot, G. B., 1966. Estuarine environmental requirements and limiting factors for striped bass. In: A Symposium on Estuarine Fisheries, Am. Fish. Soc. Special Publ. No. 3, pp. 37-49.
- Trembley, F. J., 1960. Research project on effects of condenser discharge water on aquatic life. Institute of Resrch., Lehigh Univ. Progress Report 1956 to 1959, 151 p.
- Trembley, F. J., 1961. Research project on effects of condenser discharge water on aquatic life. The Institute of Resrch., Lehigh Univ. Progress Report.
- Vernon, E. H., 1958. An examination of factors affecting the abundance of pink salmon in the Fraser river. Progr. Rep. int. Pacif. Salm. Fish. Comm., 52 p.
- Wells, H. W., M. J. Wells, and I. E. Gray., 1961. Winter fish mortality in Pamlico sound, North Carolina. Ecol., 42: 217-219.

V. AQUATIC PLANTS AND BENTHOS

General

Temperature regulates molecular movement and thus largely determines the rate of chemical reactions and consequently the rate of metabolism and activity of all organisms, both those with a relative constant temperature (homoeothermus) and those with a variable temperature that is identical with, or close to, their environment (poikilothermus). Because of its regulative capacity in determining the rate of metabolism, temperature presumably is the most important single environmental entity concerning life and life processes.

Variations in temperature of streams, lakes, estuaries, and oceans are normal events that result from climatic and geologic phenomena. The range of temperatures in waters that support some form of aquatic life other than viruses or bacteria is from -3°C (26.6°F) in super-cooled sea water to 85°C (185°F) in fresh-water thermal springs; most aquatic poikilotherms, if not all, tolerate only those temperature changes that occur within a much narrower range whether it be high, intermediate, or low on this scale of temperatures.

Within the same species the biological effects of a given temperature or temperature pattern may be different in different populations, at different ages, in different life cycle stages, or in the two sexes (Sprague, 1963), and such effects may depend on the temperature history of the individual

tested (Prosser, et al., 1952), as well as on present or past effects of other environmental factors. Many organisms experience temperature changes in their natural habitat, and these changes can be an important prerequisite for their well being and completion of their life cycles. A constant temperature of 20°C (68°F), temperatures fluctuating rapidly and irregularly between 15°C and 25°C (59 and 77°F) with an average value of 20°C (68°F), and temperatures fluctuating gradually and periodically between 15°C and 25°C (59°C and 77°F) with an average of 20°C (68°F) do not necessarily have the same biological effects. Thus, even if other prerequisites are satisfactory, the absolute temperature values of a body of water are only one measure of its suitability for a normal assemblage of aquatic life; consideration also must be given to temperature patterns or dynamics. For example, distinctions must be made between constant and fluctuating temperatures, between gradients, ranges, averages, frequency and intensity of changes, duration of a given pattern, and total summation (Kinne, 1963).

The temperature range tolerated by many species of organisms is narrow during very early development, then increases somewhat and finally decreases again in the "old adult." Similarly, it often is more restricted during the sexual phase than during other phases. Upper lethal temperatures may be lower for animals from cold water than for closely related species from warm water (Prosser and Brown, 1961). A similar aspect presumably could be found among closely related algae as well as other plants. Many mobile

organisms such as fish, some zooplankton, certain algae, and bottom-associated animals can avoid critical temperatures by vertical and horizontal migration into more suitable conditions. Other such organisms may be attracted to areas with critical temperatures and succumb when these are attained.

Fresh-Water Algae and Other Aquatic Plants

Except for fresh waters in tropical areas, relatively broad temperature ranges naturally occur seasonally and diurnally, but even in tropical waters there may be some narrow seasonal changes or major changes resulting from unusual climatic events. Although tropical waters usually experience only minor temperature changes, significant qualitative and quantitative modifications occur in their associated flora. Similar changes have been found during summer in temperate and north-temperate fresh water apparently with no attendant temperature variation, and such phenomena have led to the hypothesis that temperature, per se, bears no relationship to floristic changes in these and other waters (Blum, 1953; Pearsall, 1923; and Butcher, 1924). Nevertheless, more recent studies employing laboratory procedures wherein variables other than temperature were static have shown that temperature, per se, does have a profound influence on aquatic algae and other plants (Phinney and McIntyre, 1965; Owens and Maris, 1964), and many others provide nonlaboratory data that strongly implicate temperature as one of the causes of both qualitative and quantitative changes in aquatic flora (for example: Cairns, 1956; Wallace, 1955; Trembley, 1960; Palmer,

1965; and Patrick, 1948). Extracellular algal products, nutrients, or other factors can cause algal population changes (Mackenthun, et al., 1964; Mackenthun, 1965; Hartman, 1960; and Fogg, 1962), and these may have been responsible for those population changes mentioned above where there were no attendant temperature variations.

Algae and other plants, like poikilothermous animals, lack physiological mechanisms to maintain constant internal temperatures, and have tissue temperatures identical with or very close to that of their environment. Terrestrial plants are subjected to much wider temperature ranges than those living in aquatic environments, but a few species of aquatic algae tolerate temperatures higher than any terrestrial plants. For example, several authors have reported that some species of algae can tolerate water temperatures as high as 85°C (175°F) as found in thermal springs (Mann and Schlichting, 1967; and Kinne, 1963), but optimal temperatures for the same or similar species may range from 51° to 56°C (123.8° to 132°F) according to Brock and Brock, 1966. Other algae, notably certain diatoms, can tolerate low temperatures near 0°C (32°F), and some may remain viable after freezing. Ulothrix zonata, a filamentous green algae, grows best below 15°C (59°F), and can produce reproductive bodies, zoospores, at temperatures near 0°C (32°F) in ice water (Oltmanns, 1922-23, as cited by Blum, 1953).

Such tolerances to high or low temperature extremes are not universal among algae and other aquatic plants; similarly, an individual plant may

not thrive at all temperatures between those extremes mentioned above. Rather, there appears to be particular temperature ranges that are tolerated by each species and by closely related species or groups of species. A similar concept applies to optimum temperatures for aquatic algae. Thus, Cairns (1956) indicated that in an unpolluted stream diatoms grow best at 18° to 20°C (64.4 to 68°F); green algae at 30°C (86° to 95°F); and blue-green algae at 35° to 40°C (95 to 104°F). If environmental temperatures near 10°C (50°F) are increased either naturally or artificially to about 38° (100.4°F), the predominance of groups of species changes correspondingly from diatoms to green algae and finally at the uppermost temperatures to blue-green algae (Wallace, 1955). A few of the more high-temperature-tolerant species belonging to algal groups other than the blue-green may persist with the predominant blue-green species in such cases, and several less tolerant species of blue-greens may succumb with the diatoms and green algae at these higher temperatures.

Bottom Organisms

In a study by Strangenberg and Pawlaczyk (1961) on the effect of warm-water discharges on

a river they found that river-bottom plants and animals decreased in number when the water temperature exceeded 30°C (86°F). The macro-invertebrate riffle fauna of the Delaware River was adversely affected by heated water effluents. The macro-invertebrate biomass was reduced from 1.04 to 0.09 grams per square foot throughout the summer in the area of maximum heated water, as compared with a control station. A 35°C (95°F) water temperature at the time of sampling was found to be causing a detrimental effect on many organisms, especially the caddisfly, Hydropsyche sp., many of which were dead, while those alive were extremely sluggish. The data suggest that there is a tolerance limit close to 32.2°C (90°F) for a variety of different kinds of animals in the population structure of benthos with extensive losses in numbers and diversity of organisms accompanying further temperature increase (Coutant, 1962).

Another classic demonstration on the effects of increasing water temperatures upon the change in the composition of a macro-invertebrate population is presented by Walshe (1948). The thermal index (22-hour LD_{50}) of seven species of midge larvae reflect the probable sequence of preferred temperatures. These seven species and their thermal indices are as follows: Tanytarsus brunnipes, 29°C (84.2°F); Prodiamesa olivacea, 30°C (86°F); Anatopynia nebulosa, 30.4°C (95°F); C. longistylus, 35°C (95.9°F); and Anatopynia varia, 38°C (101.8°F).

In studies on the shift of the composition of macro-invertebrate populations (Wurtz and Renn, 1965), it was shown that no immediate kills resulted from thermal shock of 14°C (25°F). However, persistent exposure to 35°C (95°F) over 24 hours brought about changes in the composition of the macro-invertebrate population.

Studies of particular species of macro-invertebrates have shown that lethal temperatures vary considerably with the type of organism. Noland and Reichel (1943) in studying the fresh-water snail (Lymnaea stenalis) found that cultures died when the water temperature reached 30.5°C (89.6°F). Fresh-water snails (Viviparus malleatus) died when held at a temperature of 37.5°C (99.5°F) (Hutchinson, 1947).

The highest 24-hour median tolerance limit lethal temperatures that could be obtained by raising acclimation temperatures from 10°C (50°F) to 20°C (68°F) were estimated to be 34.6°C (94.2°F) for the sowbugs Asellus intermedius, Forks, and the scud, Gammarus fasciatus, Say, 33.2°C (91.8°F) for the scud, Hyallella azteca (Saussure), and 29.6°C (85.3°F) for the scud, Gammarus pseudolimnaeus Bousfield (Sprague, 1963).

The fresh-water snail (Physa gyrina) has been found to live and reproduce in a waste water ditch between 28° and 35°C (82.4° to 95°F) (Agersborg, 1932).

Many marine bottom-associated organisms have stenothermal or narrow temperature ranges. In some cases, a particular species may be stenothermal

for one developmental stage, and eurythermal for another. Breeding or spawning requirements are generally stenothermal. The time of spawning for molluscs is highly dependent upon temperature. Most molluscs with specific temperature-breeding relationships are spring and summer spawners, and many do not spawn until a certain temperature is reached (Allen, 1963). Spawning of the American oyster (Crassostrea virginica) takes place between 15 and 32 to 34°C (59 and 89.6 to 93.2°F), depending on condition of the oyster, and the spawning process is usually triggered by a rise in temperature (Galtsoff, 1964).

A large number of species are able to tolerate higher temperatures than those at which they can breed. For example, Carcinus maenas thrives, but does not breed in 14-28°C temperatures (57.2-82.4°F) (Naylor, 1965). In the case of the European lobster, temperature controls a different part of reproduction. Larvae require a minimum temperature of 15°C (59°F) even though the developing eggs, and adults, will tolerate lower temperatures (Gunter, 1957). For the above two cases, temperature limits the populations and recruitment of organisms must occur from outside the heated area.

Physiology, metabolism, and development are all affected by temperature. At a temperature of 6-7°C (42.8-44.6°F), C. virginica ceases feeding. Above 32°C (89.6°F) ciliary action, responsible for movement of water, rapidly decreases; and almost all functions of the body cease, or are reduced to a minimum at 42.0°C (107.6°F). The European

oyster, Ostrea lurida, has a tendency to close it's shell in response to falling temperatures. At 4-6°C (39.2-42.8°F) the shells of oysters remain closed most of the time; at 6-8°C (42.8-46.4°F) the shells open for about 6 hours per day; and at 15°C (59°F) the shells remain open for 23 hours per day (Galtsoff, 1964). Very little is known about prolonged effects of temperature above 32-34°C (90-94°F) on oyster populations. Long, continued exposure to high temperatures may impede the normal rate of water transport. When either low or high temperatures cause a closing of shells or a ceasing of ciliary action, oysters cease to feed and lose weight. Thus, temperature may produce an effect similar to chronic toxicity.

Acclimation and tolerance of bottom-associated organisms may be affected by temperature changes. The crab (Hemigrapsus nudus) can regain tolerance to high temperatures, after a low temperature history, in less than a week. Shore crabs (Pachygrapsus crassipes) may require a half time of six days in order to acclimate to a temperature change of 7.5°C (13.5°F) (Kinne, 1967). The giant scallop (Placopecten magellanicus) acclimates rapidly to a rise in temperature of 1.7°C (3.1°F) per day, but may take as long as three months to lose this acclimation to high temperatures (Dickie, 1958). The opossum shrimp (Neomysis americana) is very intolerant to temperature increases, and does not appear to survive at temperatures above 31°C (87.8°F) in the Chesapeake estuary (Mihursky and Kennedy, 1967).

Distribution of benthic organisms may be controlled by temperature. Reef-forming corals will not live where temperature falls below $18-19^{\circ}\text{C}$ ($64.4-66.2^{\circ}\text{F}$). The American oyster (C. virginica) on the Gulf coast is present in water that may vary from $4-34^{\circ}\text{C}$ ($39.2-93.2^{\circ}\text{F}$); while the European oyster (O. edulis) is restricted to water with temperatures of $0-20^{\circ}\text{C}$ ($32-68^{\circ}\text{F}$) (Gunter, 1957). In a study on the York River, in Virginia, Warinner and Brehmer, 1966, found that the community composition and abundance of marine benthic invertebrates in the river were affected by thermal discharge over a distance of 300-400 meters from the discharge outfall. They concluded that during the months of high normal river temperatures there was clear evidence of biological stress.

Cold is as important as heat in its effects on marine organisms. Cold water may kill directly, or in some cases indirectly when organisms are "numbed" or rendered inactive and unable to protect themselves from predators (Gunter, 1957).

One of the benefits derived from heated water is the defouling of intake pipes. Experiments have shown that fouling by Mytilus edulis and M. californianus could be controlled by tri-weekly reversals of heated discharge water either for periods of one hour at $38-41^{\circ}\text{C}$ ($100.4-105.8^{\circ}\text{F}$) or for seven hours at 34.5°C (91.4°F) (Naylor, 1965).

Summary

1. Temperature is one of the most important single environmental entities concerning life and life processes. The various functions of an organism may have somewhat different temperature ranges, and if these are not provided in the habitat the organism will die.
2. When water temperatures are increased, the predominance of groups of algal species changes correspondingly from diatom to green algae and finally at higher temperatures to blue-green algae.
3. The number and distribution of bottom organisms decrease as water temperatures increase with a tolerance limit close to 90°F for a "balanced population structure. Studies of particular species of macro-invertebrates have shown that lethal temperatures vary considerably with the type of organism. In some cases a particular species may be stenothermal for one developmental stage, and eurythermal for another. Thus, a large number of species are able to tolerate higher temperatures than those at which they can breed.
4. Cold is as important to aquatic plants and benthos as is heat.
5. One of the beneficial uses of heated effluents is the defouling of intake pipes; accomplished by reversing the flow of water through the pipes for a specified period of time.

REFERENCES CITED

- Agersborg, H. P. K., 1932. The relation of temperature to continuous reproduction in the pulmonate snail. *Nautilus*, 45(4): 121-123.
- Allen, J. A., 1963. Ecology and functional morphology of molluscs. *Oceano. Mar. Bio. Ann. Rev.*, 1, Harold Barnes, Ed., 253-288.
- Blum, J. L., 1953. The ecology of algae growing in the Saline River, Michigan, with special reference to water pollution. Doc. Thesis, Univ. Mich., ix + 176 pp.
- Blum, J. L., 1956. The ecology of river algae. *Bot. Rev.*, 22: 291-341.
- Brock, T. D., and M. L. Brock, 1966. Temperature optima for algal development in Yellowstone and Iceland hot springs. *Nature*, 209, No. 5024, 733-34.
- Butcher, R. W., 1924. The plankton of the River Wharfe (Yorkshire). *Naturalist*, pp. 175-180, 211-214.
- Cairns, J., Jr., 1956. Effects of increased temperatures on aquatic organisms. *Ind. Wastes*, 1(4): 150-152.
- Coutant, C. C., 1962. The effect of a heated water effluent upon the macro-invertebrate riffle fauna of the Delaware River. *Penn. Acad. Sci.*, 37: 58-71.
- Dickie, L. M., 1958. Effects of high temperatures on survival of the giant scallop. *Jour. Fish Resch. Bd. Can.*, 15(6): 1189-1211.
- Fogg, G. E., 1962. The importance of extra-cellular products of algae in the aquatic environment. *Bio. Prob. Wat. Poll.*, 3rd Seminar, PHS Publ. No. 999-WP-25, C. M. Tarzwell, Ed., 424 pp.
- Galtsoff, P. S., 1964. The American oyster Crassostrea virginica Gmelin. U. S. Fish & Wildlife Serv., Spec. Sci. Reps., Fish. No. 64, 480 p.
- Gunter, G., 1957. Temperature. *Treatise on marine ecology and palaeoecology I.* (Hedgepeth, J. W., Ed.), Chap. 8, Geol. Soc. Am. Mem., 67, 159-184.
- Hartman, R. T., 1960. Algae and metabolites of natural waters. In: *The Ecology of Algae*, Spec. Publ. No. 2, Pymatuning Lab. of Field Biology, Univ. Pittsburgh, Pittsburgh, Pennsylvania, pp. 38-55.
- Hutchinson, L., 1947. Analysis of the activity of the fresh water snail, Viviparus malleatus (Reeve). *Ecol.*, 28(4): 335-345.

Kinne, O., 1963. The effect of temperature and salinity on marine and brackish water animals. Temperature. In: Oceano. and Mar. Bio., Ann. Rev., 1 (Harold Barnes, Ed.), 301-340.

Kinne, O., 1967. Physiology of estuarine organisms with special reference to salinity and temperature: general aspects. In: Estuaries, George H. Lauff, Ed., Amer. Assoc. Adv. Sci., Publ. No. 83, 525-540.

Mackenthun, K. M., W. M. Ingram, and R. Porges. 1964. Limnological Aspects of Recreational Lakes. U. S. Public Health Service, Publ. No. 1167: 176 pp.

Mackenthun, K. M.

1965. Nitrogen and phosphorus in water, an annotated selected bibliography of their biological effects. U. S. Public Health Service, Publ. No. 1305: 111 pp.

Mann, J. E. and H. E. Schlichting, Jr., 1967. Benthic algae of selected thermal springs in Yellowstone National Park. Trans. Amer. Microscopical Soc., 86(1): 2-9.

Mihursky, J. A., and V. S. Kennedy, 1967. Water temperature criteria to protect aquatic life. Symposium on Water Quality Criteria to Protect Aquatic Life, Am. Fish. Soc., Spec. Publ. No. 4, 20-32.

Naylor, E., 1965. Biological effects of a heated effluent in docks at Swansea, S. Wales. Proc. Zool. Soc., London, 144: 253-268; Water Poll. Abstr., 39(6): Abstr. No. 997.

Naylor, E., 1965. Effects of heated effluents upon marine and estuarine organisms. In: Advances in Marine Biology, 3, Academic Press, New York, 63-103.

Noland, L. E., and E. Reichel, 1943. Life cycle of Lymnaea stagnalis completed at room temperature without access to air. Nautilus, 57(1): 8-13.

Oltmanns, F., 1922-23. Morphologic und biologic der algen. 2 auff., 3 vols.

Owens, M., and P. J. Maris, 1964. Some factors affecting the respiration of some aquatic plants. Hydrobiologia, 23: 533-543.

Palmer, C. M., 1965. Phytoplankton periodicity in a Newfoundland pond. Phycologia, 1: 3940.

Patrick, R., 1948. Factors affecting the distribution of diatoms. Bot. Rev., 14(8): 478-524.

Pearsall, W. H., 1923. A theory of diatom periodicity. Jour. Ecol., 11: 165-183.

- Phinney, H. K., and C. D. McIntire, 1965. Effect of temperature on metabolism of periphyton communities developed in laboratory streams. *Limnol. Oceanog.*, 10(3): 341-344.
- Prosser, C. L., et al., 1952. Comparative animal physiology. W. B. Saunders Co., Philadelphia, Pa., 888 pp.
- Prosser, C. L., and F.A. Brown, Jr., 1961. Comparative animal physiology. Second Edition, W. B. Saunders Co., Philadelphia, Pa., 688 pp.
- Sprague, J. B., 1963. Resistance of four freshwater crustaceans to lethal high temperature and low oxygen. *Jour. Fish. Resch. Bd. Can.*, 20(2): 387-415.
- Trembley, F. J., 1960. Research project on effects of condenser discharge water on aquatic life, progress report, 1956 to 1959. Institute of Resch. Lehigh Univ., 154 pp.; *Water Poll. Abstr.*, 34(11), Abstr. No. 2157.
- Wallace, N. W., 1955. The effect of temperature on the growth of some freshwater diatoms. *Notulae naturae of the Acad. Nat. Sci. of Philadelphia*, 280: 1-11.
- Walshe, B. M., 1948. Oxygen requirements and thermal resistance of chironomid larvae from flowing and still water. *Jour. Exp. Bio.*, 25: 35.
- Warinner, J. E., and M. L. Brehmer, 1966. The effects of thermal effluents on marine organisms. *International Jour. Air and Water Poll.*, 10(4): 277-289.
- Wurtz, C. B., and C.E. Renn, 1965. Water temperatures and aquatic life. Prepared for Edison Electric Institute Research Project No. 49, 99 pp.

TEMPERATURE and AQUATIC LIFE

A Selected Bibliography

- Abbott, B. J., 1960. A note on the oxygen and temperature tolerance of the triclads, Phagocata gracilis (Haldeman) and Dugesia tigrina (Girard). Virginia Jour. Sci., 2(1): 1-8.
- Adlung, K. G., 1957. The toxicity of insecticides to fish and its dependence on temperature. Nat. Wiss., 44: 622-623. Water Poll. Abstr., 31(12), Abstr., No. 2446, 1958. Spo. Fish. Abstr., 4(1), Abstr. No. 2166, 1959.
- Agersborg, H. P. K., 1930. The influence of temperature on fish. Ecology, 11(1): 136-144.
- Agersborg, H. P. K., 1932. The relation of temperature to continuous reproduction in the pulmonate snail. Nautilus, 45(4); 121-123.
- Alabaster, J. S., 1962. The effect of heated effluents on fish. Internat. Conf. on Water Poll. Resrch., London, 1962. Air and Water Poll. 7(6/7): 541-563, 1963. Bio. Abstr., 45(13), Abstr. No. 54305, 1964.
- Alabaster, J. S. and A. L. Downing, 1960. The behavior of roach (Rutilus rutilus L.) in temperature gradients in large outdoor tanks. Proc. Indo-Pacific Fish. Coun. III(49).
- Alabaster, J. S. and A. L. Downing, 1966. A field and laboratory investigation of the effect of heated effluents on fish. Fish. Invest. (Minist. of Agr., Fish. and Food of Unit. King.), Ser. I, 6(4): 42.
- Alabaster, J. S., D. W. M. Herbert, and J. Hemens, 1957. The survival of rainbow trout (Salmo gairdneri Richardson) and perch (Perca fluviatilis L.) at various concentrations of dissolved oxygen and carbon dioxide. Ann. Appl. Bio. 45: 177-188.
- Alabaster, J. S., and K. G. Robertson, 1961. Effect of diurnal changes in temperature, dissolved oxygen, and illumination on the behavior of roach (Rutilus rutilus L.), bream (Abramis brama L.) and perch (Perca fluviatilis L.). Animal Behavior, 9(3-4): 187.
- Alabaster, J. S., and A. Swain, 1963. Heated water and fish. Ann. Rep. Challenger Soc., 3(15): 39.
- Alabaster, J. S., and R. L. Welcomme, 1962. Effect of concentration of dissolved oxygen on survival of trout and roach in lethal temperatures. Nature, London, 194: 107. Water Poll. Abstr., 36: 672, 1963. Spo. Fish. Abstr., 9(2), Abstr. No. 6587, 1964.

Alderdice, D. F., 1963. Some effects of simultaneous variation in salinity, temperature and dissolved oxygen on the resistance of young coho salmon to a toxic substance. Jour. Fish. Res. Bd. Can., 20: 525-550. Spo. Fish. Abstr. 8(3), Abstr. No. 5905, 1963. Water Poll. Abstr. 37(1), Abstr. No. 178, 1964.

Allanson, B. R., and R. G. Noble, 1964. The tolerance of Tilapia mossambica (Peters) to high temperature. Trans. Am. Fish. Soc. 93(4): 323-332. Spo. Fish. Abstr. 10(1), Abstr. No. 7192, 1965.

Allbaugh, C. A., and J. V. Manz., 1964. Preliminary study of the effects of temperature fluctuations on developing walleye eggs and fry. Prog. Fish-Cult. 26(4): 175-180.

Allee, W. C., A. E. Emerson, O. Park, T. Park, and K. P. Schmidt, 1949. Principles of animal ecology. W. B. Saunders Co., Philadelphia, Pa.

Allen, K. R., 1940. Studies on the biology of the early stages of the salmon, (Salmo salar): I. Growth in the River Eden. Jour. Animal Ecol. 9(1): 1-23.

Ames, A. M., and W. W. Smith, 1944. The temperature coefficient of the bactericidal action of chlorine. Jour. of Bact., 47: 445. Water Poll. Abstr., 17, Jul. 1944.

Anaichev, A. V., 1959. Digestive enzymes of fish and seasonal changes in their activity. Biokhimiia (Transl.), 24(6): 952-958. Spo. Fish. Abstr., 6(1), 1960.

Anderson, R. C., 1951. Preferred temperature of a sample of Lepomis gibbosus, the pumpkinseed. Manu. in the Ontario Fish. Resrch. Lab. Library, Toronto.

Andrews, C. W., 1946. Effect of heat on the light behavior of fish. Trans. Roy. Soc. of Can., Ser. 3, 40, Soc. 5: 27-31.

Angelovic, J. W., W. F. Sigler, and J. M. Neuhold. Temperature and flourosis in rainbow trout. Jour. Water Poll. Control Fed., 33: 371-381, Apr. 1961.

Anon., 1949. Discharge of heated liquids into streams. Rep. of Rivers Poll. Prevention Sub-Committee of the Central Water Committee, Minist. of Health, England, pp: 69-76.

Anon., 1954. Changes in the dissolved-oxygen content of river water used for direct cooling. Rep. No. 10, Generating Station Operation-Resrch. Liaison Committee. Brit. Electr. Author.

Anon., 1955. First progress report. Aquatic Life Advisory Committee of ORSANCO, Sewage and Industrial Wastes, 27(3): 321-331.

- Anon., 1956. Feed heavy on a rise in temperature. Prog. Fish Cult., 18(1): 44.
- Anon., 1956. Proceedings of the fifth annual water symposium, February 1956. Louisiana State Univ., Engng. Exp. Sta. Bull. 55.
- Anon., 1956. Aquatic life water quality criteria (Second progress report of the Aquatic life Advisory Committee of the Ohio River Valley Water Sanitation Commission). Sewage and Indus. Wastes, 28(5): 678-690.
- Anon., 1958. Oxygen relationships in streams, proc. of seminar at Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio, Oct. 30 - Nov. 1, 1957. Publ. Health Serv. Tech. Rep. No. W58-2.
- Anon., 1960. Is heat a pollutant? Pennsylvania thinks so. Chem. Engng. Prog. Staff, Chem. Engng. Prog. 56: 33.
- Anon., 1961. Effect of water temperature on stream re-aeration. Proc. Am. Soc. Civ. Engrs. 87, SA6: 59-71.
- Anon., 1962. Heated discharges . . . their effect on streams. Rep. by the Advisory Committee for the control of stream temperatures to the Pa. Sanitary Water Board. Pa. Dept. Health, Harrisburg, Publ. No. 3, 108 pp.
- Anon., 1963. Water temperature influences, effects, and control. Proc. of the 12th Pacific Northwest Symposium on Water Poll. Resrch., Pacific Northwest Water Lab., Publ. Health Serv., U. S. Dept. of Health, Education and Welfare, 160 pp.
- Anon., 1963. Sediment transportation mechanics: density currents. Prog. Rep., Task Committee on Preparation of Sedimentation Manual, Committee on Sedimentation, Jour. of the Hydraulics Div., ASCE, 89(HY5), Proc. Paper 3639, pp. 77-87.
- Anon., 1965. Symposium on streamflow regulation for quality control. U. S. Publ. Health Serv. Publ. No. 999-WP-30, 420 pp.
- Anon., 1965. Minutes, third conference on Patuxent estuary studies, Nov. 13-14, 1964. Chesapeake Bio. Lab. Ref. No. 65-23.
- Anon., 1966. Nuclear power plant proposal raises pollution issues. Water Control News, 1(1), May 16.
- Anon., 1967. Temperatures for hatching walleye eggs. Prog. Fish Cult., 29(1): 20 (Jan.).
- Ansell, A. D., 1963. Venus mercenaria L. in Southampton water. Ecology 44(2): 396-397.

Ansell, A. D., 1963. The biology of Venus mercenaria in British waters, and in relation to generating station effluents. Ann. Rep. Challenger Soc. 3(15): 38.

Ansell, A. D., and F. A. Loosemore, 1963. Preliminary observations on the relation between growth, spawning and condition in experimental colonies of Venus mercenaria L. Jour. du Conseil, 28: 285-294.

Ansell, A. D., K. F. Lander, J. Coughlan, and F. A. Loosemore, 1964. Studies on the hard-shell clam, Venus mercenaria, in British waters. I Growth and reproduction in natural and experimental colonies. Jour. Appl. Ecol., 1: 63-82.

Ansell, A. D., F. A. Loosemore, and K. F. Lauder, 1964. Studies on the hard-shell clam, Venus mercenaria, in British waters. II Seasonal cycle in bio-chemical composition. Jour. Appl. Ecol., 1: 83-95.

Anthony, E. H., 1961. The oxygen capacity of goldfish blood (Carassius auratus) in relation to thermal environment. Jour. Exp. Bio. 38: 93-107.

Armitage, K. B., 1962. Temperature and O₂ consumption of Orchomonella chilensis (Heller) (Amphipoda: Gammaroida). Bio. Bull. 123(2): 225-232.

Arnold, G. E., 1962. Thermal pollution of surface supplies. Jour. Am. Water Works Asso., 54(11): 1332-1346.

Bailey, R. M., 1955. Differential mortality from high temperature in a mixed population of fishes in southern Michigan. Ecol., 36(3): 526-528. Spo. Fish. Abstr. 1(4), Abstr. No. 519, 1956.

Bakastov, S. S., 1960. Some data on bottom temperatures in the Rybinsk Reservoir when the surface is frozen. Bull. Inst. Bio. Vodokhranilishch Akad. Nauk SSSR 8/9: 62-66. Spo. Fish. Abstr. 7(4), 1962.

Bakshtanskii, E. L., 1961. The role of feeding and of warming of the water in the artificial rearing of salmon above the Arctic circle. Rybnoe Khoz. 10: 15-18. Referat. Zhur., Bio. 1962, No. 9178. Spo. Fish. Abstr. 8(2), 1963.

Baldwin, E., 1948. An introduction to comparative biochemistry. Cambridge Univ. Press, 164 pp.

Baldwin, N. S., 1957. Food consumption and growth of brook trout at different temperatures. Trans. Am. Fish. Soc. 86: 323-328. Spo. Fish. Abstr. 3(1), Abstr. No. 1362, 1958.

Banta, A. M., and T. R. Wood, 1928. A thermal race of cladocera originated by mutation. Zeitschr. Induct. Abstam - u. Vererbungsl. Supplement b, 1: 397-398.

Baranov, I. V., 1961. Thermic and hydrochemical condition in the Gorikovskii Reservoir. Trudy Inst. Bio. Vodokhranilishch Akad. Nauk SSSR 4(7): 294-320. Referat. Zhur., Bio., 1962, No. 11Zh332. Spo. Fish. Abstr., 8(2), 1963.

- Bardach, J. E., 1955. Certain biological effects of thermocline shifts. *Hydrobiologia*, 7(4): 309-324.
- Bardach, J. E., and R. G. Bjorklund, 1957. The temperature sensitivity of some American freshwater fishes. *Am. Nat.* 91(859): 233-251. *Spo. Fish. Abstr.* 3(1), Abstr. No. 1322, 1958.
- Barges, H. M., 1950. Fish distribution studies, Niaugua Arm of the Lake of the Ozarks, Missouri. *Jour. Wildl. Mgnt.* 19(1): 16-33.
- Barrington, E. J. W., and A. J. Matty, 1954. Seasonal variation in the thyroid gland of the minnow, Phoxinus phoxinus L. with some observations on the effect of temperature. *Proc. Zool. Soc. of London*, 124: 547-564.
- Basu, S. P., 1959. Active respiration of fish in relation to ambient concentrations of oxygen and carbon dioxide. *Jour. Fish. Resrch. Bd. Can.* 16(2): 175-212.
- Bata, G. L., 1957. Recirculation of cooling waters in rivers and canals. *Jour. of the Hydraulics Div. ASCE*, 83(HY3), Proc. Paper 1265.
- Battle, H. I., 1926. Effects of extreme temperature on muscle and nerve tissue in marine fishes. *Trans. Proc. and Roy. Soc. Can.*, 5(20): 127-143.
- Battle, H. I., 1929. Effects of extreme temperatures and salinities on the development of Enchelyopus cimbrius L. *Contr. Can. Bio. (N.S.)* 5: 109-192.
- Baudin, L., 1926. Variation des echanges respiratoires des poissons en fonction des la pression atmospheriques et de la temperature. *Mem. Soc. Vaudoise, Sci. Nat.* 4(1): 1-40.
- Beamish, F. W. H., 1964. Respiration of fishes with special emphasis on standard oxygen consumption. II. Influence of weight and temperature on respiration of several species. *Can. Jour. Zool.* 42: 161-188. *Water Poll. Abstr.* 37(10), Abstr. No. 1564, 1964.
- Beaven, G. F., 1960. Temperature and salinity of surface water at Solomons, Maryland. *Chesapeake Sci.*, 1(1): 2-11.
- Becker, H. G., 1924. Mechanism of absorption of moderately soluble gases in water, and its use in determining the rate of solution of oxygen by water under different conditions of mixing. *Indus. and Engng. Chem.*, 16(12): 1220-1224.
- Beckman, W. C., 1942. Annulus formation on the scales of certain Michigan game fishes. *Papers of the Michigan Acad. of Sci., Arts and Letters* 28:281-312.
- Beeton, A., 1958. The vertical migration of Mysis relicta in Lakes Huron and Michigan. *Dissertation Abstr.*, 19(3): 601. *Spo. Fish. Abstr.*, 3(3), 1958.

- Beeton, A. M., 1960. Environmental changes in Lake Erie. Trans. Am. Fish. Soc., 90(2): 153-159.
- Belchradek, J., 1930. Temperature coefficients in biology. Bio. Review, 5(1): 30-58.
- Belding, D. L., 1928. Water temperature and fish life. Trans. Am. Fish. Soc., 58: 98-105.
- Belehradek, J., 1931. Influence de la temperature sur la frequence cardiaque chez les embryons de la rousette, Scylliorhinus canicula L. C. R. Soc., Bio., Paris 107(20): 727-729.
- Belehradek, J., 1935. Temperature and living matter. Protoplasmanonographien 8, Berlin, 229 pp.
- Beliaev, E. V., 1958. Experiments on raising Daphnia magna during the winter. M. Rybnaia Promyshlennost, 34: 52-55. Spo. Fish. Abstr, 5(1), 1960.
- Bell, H. S., 1942. Stratified flow in reservoirs and its prevention in silting. Misc. Publ. 491, U. S. Dept. of Agr.
- Bennett, G. W., 1965. The environmental requirements of centrarchids with special reference to largemouth bass, smallmouth bass and spotted bass. Bio. Problems in Water Poll., Third Seminar, 1962 (Tarzwell, C. M., ed.), Robert A. Taft Engng. Center, Publ. Health Serv. Publ. No. 999-WP-25, pp. 156-159.
- Bennett, G. W., and W. F. Childers, 1957. The smallmouth bass, Micropterus dolomieu, in warm water ponds. Jour. Wildl. Mgmt., 21(4): 414-424.
- Berg, K., 1943. Physiographical studies on the river Susaa. Folia Limnol. Scand., No. 1, 174 pp.
- Berg, K., 1952. On the O₂ consumption of Ancyliidae (Gastropoda) from an ecological point of view. Hydrobiologia, 4(3): 225-267.
- Berg, K., 1953. The problem of respiratory acclimatization. Hydrobiologia 5: 331-350.
- Berger, B. B., 1961. Does production of power pollute our rivers? Power Engng. (Mar.).
- Berry, J. W., and P. C. Holt, 1959. Reactions of two species of Branchiobdellidae (Oligochaeta) to high temperatures and low oxygen tensions. Tech. Bull. Va. Agr. Exp. Sta., 141, 12 pp. Sewage and Indus. Wastes, 34(7), 1962.

- Biever, K. D., and M. S. Mulla, 1966. Effects of temperature on the developmental stages of Psychoda alternata (Diptera Psychodidae). Mosquito News, 26(3): 416-419.
- Binet, L., and A. Marin, 1934. Action de la chaleur sur les poissons Jour. Physiol Et. Path. Gen. 32(2): 372-379.
- Binet, L., and A. Marin, 1936. Contributions a l'etude de l'hypertermie et de l'asphyxie (recherches sur les poissons). Bio. Med., Paris, 26: 329-361.
- Birge, E. A., C. Juday, and H. W. March, 1927. The temperature of the bottom deposits of Lake Mendota: a chapter in the heat exchanges of the lake. Trans. Wisc. Acad. Sci., Arts and Letters 23: 187-231.
- Bishai, H. M., 1960. Upper lethal temperatures for larval salmonids. Jour. Cons. Int. Explor. Mer., 25: 129-133. Spo. Fish. Abstr., 5(4), Abstr. No. 3803, 1960.
- Bisset, K. A., 1946. The effect of temperature on non-specific infections of fish. Jour. Pathol. and Bactol. 58(2): 251-258.
- Bisset, K. A., 1948. Seasonal changes in the normal flora of fresh water fish. Jour. Hyg., 46(1): 94-97.
- Bisset, K. A., 1948. The effect of temperature upon antibody production in cold-blooded vertebrates. Jour. Pathol. and Bactol., 60(1): 87-92.
- Black, E. C., 1947. Respiratory characteristics of the blood of the Atlantic salmon, Salmo salar salar L., acclimated to summer temperatures in fresh water. Trans. Roy. Soc. Can. (3) 41 (Sec. I): 198.
- Black, E. C., 1953. Upper lethal temperatures of some British Columbia freshwater fishes. Jour. Fish. Resrch. Bd. Can., 10: 196-210.
- Black, E. C., 1955. Blood levels of hemoglobin and lactic acid in some freshwater fishes following exercise. Jour. Fish. Resrch. Bd. Can. 12(6): 917-929.
- Black, E. C., 1958. Hyperactivity as a lethal factor in fish. Jour. Fish. Resrch. Bd. Can., 15(4): 573-586.
- Black, V. S., F. E. J. Fry, and E. C. Black, 1947. The influence of temperature on the respiratory tolerance of young goldfish. Proc. Nova Scotia Inst. Sci., 121: 659.
- Blaxter, J. H. S., 1957. Herring rearing, III -- The effect of temperature and other factors on myotome counts. Scot. Home Dept., Mar. Res. No. 1, 16 pp.

Blaxter, J. H. S., 1960. The effect of extremes of temperature on herring larvae. Jour. Mar. Bio. Assoc. Unit. King., 39, 605.

Boetius, I., 1962. Temperature and growth in a population of Mytilus edulis L. from the Northern Harbor of Copenhagen (the Sound). Medd. fra Dan. Fisk. og Havund., N.S. 3:339-346.

Bogatova, I. B., 1962. Lethal ranges of oxygen content, temperatures, and pH for some representatives of the family Chydoridae. Zool. Zhur., 41(1): 58-62. Bio. Abstr., 40(5), Abstr. No. 21287, 1963.

Bonnet, D. D., 1939. Mortality of the cod egg in relation to temperature. Bio. Bull., Wood's Hole 76:428-441.

Borodin, N. A., 1934. Survival of fish in freezing temperatures. Jhrb., Abt. Allg. Zool. and Physiol., 53: 313-342.

Bovee, E. C., 1949. Studies on the thermal death of Hyaella azteca Saussure. Bio. Bull., 96: 123-128.

Bowen, E. S., 1932. Further studies in the aggregating behavior of Amieurus melas. Bio. Bull., 63: 258-270.

Bowen, I. S., 1926. The ratio of heat losses by conduction and by evaporation from any water surface. Physical Review, 27(2).

Boycott, A. E., 1936. The habitats of fresh water molluscs in Britain. Jour. Animal Ecol., 5: 116-186.

Brawn, V. M., 1960. Temperature tolerance of unacclimated herring (Clupea harengus L.). Jour. Fish. Resrch. Bd. Can., 17(5): 721-723.

Breder, C. M., Jr., 1927. On the temperature - oxygen toleration of brook trout. Copeia, 163: 36-39.

Breder, C. M., Jr., 1951. Studies on the structure of the fish school. Bull. Am. Museum Nat. Hist., 98: 5-27.

Breder, C. M., and R. F. Nigrelli, 1935. The influence of temperature and other factors on the winter aggregation of the sunfish, Lepomis auritus, with critical remarks on the social behavior of fishes. Ecol., 16(1): 33-47.

Brett, J. R., 1944. Some lethal temperature relations of Algonquin Park fishes. Univ. Toronto Stud. Bio. Ser. 52, Publ. Ontario Fish. Resrch. Lab., 63: 1-49.

Brett, J. R., 1946. Rate of gain of heat-tolerance in goldfish (Carassius auratus). Univ. Toronto Stud. Bio. Ser. 53, Publ. Ontario Fish. Resrch. Lab., 64:9-28.

Brett, J. R., 1952. Temperature tolerance in young pacific salmon, genus Oncorhynchus. Jour. Fish. Resrch. Bd. Can., 9(6): 265-323.

Brett, J. R., 1956. Some principles in the thermal requirements of fishes. Quarterly Rev. of Bio., 31(2): 75-87. Spo. Fish. Abstr., 2(3), Abstr. No. 1046, 1957.

Brett, J. R., 1958. Some principles in the thermal requirements of fishes. Quarterly Rev. of Bio., 31(2): 75. Ontario Fish. Resrch. Lab.

Brett, J. R., 1958. Implications and assessments of environmental stress. The Investigations of Fish-Power Problems (Larkin, P. A., ed.). H. R. MacMillan Lectures in Fisheries, Univ. British Columbia, pp. 69-83.

Brett, J. R., 1960. Thermal requirements of fish - three decades of study, 1940-1970. Bio. Problems in Water Poll., Trans. 1959 Seminar, Robert A. Taft Sanitary Engng. Center Tech. Rep. W60-3, Cincinnati, pp. 110-117.

Brett, J. R., and D. F. Alderdice, 1958. The resistance of cultured young chum and sockeye salmon to temperatures below 0°C. Jour. Fish. Resrch. Bd. Can., 15(5): 805-813. Spo. Fish. Abstr., 4(1), Abstr. No. 2252, 1959.

Brett, J. R., M. Hollands, and D. F. Alderdice, 1958. The effect of temperature on the cruising speed of young sockeye and coho salmon. Jour. Fish. Resrch. Bd. Can., 15(4): 587-605. Spo. Fish. Abstr., 4(1), Abstr. No. 2274, 1959.

Bridges, W. R., 1965. Biological problems in water pollution: effects of time and temperature on the toxicity of heptachlor and kepone to redear sunfish. Third Seminar: 1962, Publ. No. 99-WP-25, Publ. Health Serv., U. S. Dept. of Health, Education and Welfare, Washington, D. C.

Britt, N. W., 1955. New methods of collecting bottom fauna from shoals of rubble bottoms of lakes and streams. Ecol., 36(3): 524-525.

Britton, S. W., 1924. The effects of extreme temperature on fishes. Am. Jour. Physiol., 67: 411-421.

Brock, T. D., and M. L. Brock, 1966. Temperature optima for algal development in Yellowstone and Iceland hot springs. Nature, 209(5024): 733-734.

Broekhuysen, G. J., 1940. A preliminary investigation of the importance of desiccation, temperature and salinity as factors controlling the vertical distribution of certain intertidal gastropods in False Bay, South Africa. Trans. Roy. Soc. South Africa, 28: 225-292.

- Brown, L. A., 1929. The natural history of cladocerans in relation to temperature. I. Distribution and the temperature limits for vital activities. *Am. Nat.*, 63: 248-264.
- Brown, M. E., 1946. The growth of brown trout (Salmo trutta L.) III The effect of temperature on the growth of two-year-old trout. *Jour. Exp. Bio.* 22(3-4), Sec. III: 145-155.
- Brown, M. E., 1951. The growth of brown trout (Salmo trutta L.) IV The effect of food and temperature on the survival and growth of fry. *Jour. Exp. Bio.*, 28: 473-491.
- Brues, C. T., 1927. Animal life in hot springs. *Quarterly Rev. of Bio.*, 2(2): 181-203.
- Brues, C. T., 1928. Studies on the fauna of hot springs in the western United States and the biology of thermophilous animals. *Proc. Am. Acad. Arts and Sci.*, 63(4): 139-228.
- Brues, C. T., 1939. Studies on the fauna of some thermal springs in the Dutch East Indies. *Proc. Am. Acad. Arts and Sci.*, 73(4): 71-95.
- Bull, H. Q., 1936. Studies on conditioned responses in fishes. VII Temperature perception in teleosts. *Jour. Man. Bio. Assm. Unit. King.*, 21: 1-27.
- Bullock, T. H., 1955. Compensation for temperature in the metabolism and activity of poikilotherms. *Bio. Rev.*, 30(3): 311-342. *Spo. Fish. Abstr.*, 2(2), Abstr. No. 828, 1957.
- Bullough, W. S., 1939. A study of the reproductive cycle in the minnow in relation to the environment. *Proc. Zool. Soc. London*, 109: 79-102.
- Burdick, G. E., M. Lipschuetz, H. J. Dean, and E. J. Harris, 1954. Lethal oxygen concentrations for trout and smallmouth bass. *New York Fish and Game Jour.*, 1: 84. *Sewage and Indust. Wastes*, 27(6), 1955.
- Burger, J. W., 1939. Some experiments on the relation of the external environment to the spermatogenic cycle of Fundulus heteroclitus L. *Bio. Bull.*, 77: 96-103.
- Burnson, B., 1938. Seasonal temperature variations in relation to water treatment. *Jour. Am. Water Works Asso.*, 30: 793.
- Burrows, R. E., 1963. Water temperature requirements for maximum productivity of salmon. Water temperature influences, effects, and control. *Publ. Health Serv., Dept. of Health, Education and Welfare, Pacific Northwest Water Lab., Corvallis, Oregon*, pp. 29-38.
- Burton, G. W., and E. P. Odum, 1945. The distribution of stream fish in the vicinity of Mountain Lake, Virginia. *Ecol.*, 26(2): 182-194.

- Butler, P. A., 1965. Biological problems in water pollution: reaction of some estuarine mollusks to environmental factors. Third Seminar: 1962, Publ. No. 999-WP-25, Publ. Health Serv., U. S. Dept. of Health, Education and Welfare, Washington, D.C.
- Butterfield, C. T., and E. Wattie, 1946. Influence of pH and temperature on the survival of coliform and enteric pathogens when exposed to chloramine. Publ. Health Rep. 61: 157, Reprint No. 2692.
- Butterfield, C. T., E. Wattie, S. Megregian, and C. W. Chambers, 1943. Influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to free chlorine. Publ. Health Rep., 58, Dec. 17, 1943.
- Byrd, I. B., 1951. Depth distribution of the bluegill, Lepomis macrochirus Rafinesque, in farm ponds during summer stratification. Trans. Am. Fish. Soc., 81:162.
- Cadwallader, L. W., 1964. Thermal pollution of watercourses. Proc. 19th Indus. Waste Conf., Purdue Univ., Engng. Extn. Serv., 117: 9-11. Water Poll. Abstr., 39(4), Abstr. No. 470, 1966.
- Cairns, J., Jr., 1956. Effects of increased temperature on aquatic organisms. Indus. Wastes, 1(4): 150-152.
- Cairns, J., Jr., 1956. Effect of heat on fish. Indus. Wastes, 1(5): 180-183.
- Cairns, J., Jr., and A. Scheier, 1958. The effects of temperature and hardness of water upon the toxicity of zinc to the pond snail, Physa heterostropha (Say). Notulae Naturae of the Acad. of Nat. Sci. of Phila., 308.
- Cairns, J., Jr., and A. Scheier, 1964. The effects of sublethal levels of zinc and of high temperature upon the toxicity of a detergent to the sunfish Lepomis gibbosus L. Notulae Naturae, 367: 1-3. Bio. Abstr., 45, Abstr. No. 94311, 1964.
- Camp, T. R., D. A. Root, and B. V. Bhoota, 1940. Effects of temperature on rate of floc formation. Jour. Am. Water Works Asso., 32: 1913.
- Carlisle, D. B., 1957. On the hormonal inhibition of moulting in decapod crustacea. II The terminal anecdysis in crabs. Jour. Mar. Bio. Asso. Unit. King., 36: 291-307.
- Chadwick, W. L., F. S. Clark, and D. L. Fox, 1950. Thermal control of marine fouling at Redondo Steam Station of the Southern California Edison Company. Trans. ASME, Feb.
- Chalkley, H. W., 1930. High lethal temperature, paramecium. Physiol. Zool., 3: 425-440.

Chang, S. L., M. Buckingham, and M. P. Taylor, 1948. Studies on Leptospira icterohaemorrhagiae, IV Survival in water and sewage, destruction in water by halogen compounds, synthetic detergents and heat. Jour. of Infectious Diseases, 82: 256.

Chang, S. L., and G. M. Fair, 1941. Viability and destruction of cysts of Endameba histolytica. Jour. Am. Water Works Asso., 33: 1708-1709.

Chellis, R. D. and E. Ireland, 1959. Site studies for a steam power plant. Jour. of the Power Div., ASCE, 85(P01), Proc. Paper 1949, Feb.

Chidester, F. E., 1924. A critical examination of the evidence for physical and chemical influences on fish migration. Brit. Jour. Exp. Bio., 2: 79-118.

Churchill, M. A., 1947. Effect of density currents upon raw water quality. Jour. Am. Water Works Asso., 39(3): 357.

Churchill, M. A., 1945. Discussion of "translatory waves." Trans. ASCE 110: 1229.

Churchill, M. A., 1954. Analysis of a stream's capacity of assimilating pollution. Sewage and Indus. Wastes, 26(7).

Churchill, M. A., 1958. Effects of impoundments on oxygen resources. Oxygen Relationships in Streams, Proc. of Seminar, Cincinnati, Ohio, Oct. 30-Nov. 1, 1957. Tech. Rep. W58-2: 107-124.

Churchill, M. A., 1958. Effects of storage impoundments on water quality. Trans. ASCE, 123: 419.

Churchill, M. A., 1965. Control of temperature through streamflow regulation. Streamflow Regulation for Quality Control, Proc. of Symposium, Cincinnati, Ohio, Apr. 3-5, 1963, Publ. Health Serv. Publ. No. 999-WP-30: 179-192.

Clark, D., 1959. River inadequate for cooling needs of new power stations. Surveyor, London, 118: 689-690.

Clausen, R. G., 1931. Orientation in fresh water fishes. Ecol., 12(3): 541-546.

Clausen, R. G., 1933. Fish metabolism under increasing temperature. Trans. Am. Fish. Soc., 63: 215-219.

Clausen, R. G., 1939. Body temperature of fresh water fishes. Ecol., 15(2): 139-144.

- Clemens, H. P., and K. E. Sneed, 1958. Effect of temperature and physiological condition on tolerance of channel catfish to pyridylmercuric acetate (PMA). *Prog. Fish. Cult.*, 20(4): 147-150.
- Cocking, A. W., 1957. Relation between the ultimate upper lethal temperature and the temperature-range for good health in the roach (Rutilus rutilus). *Nature*, 180: 661-663.
- Cocking, A. W., 1959. The effects of high temperature on roach (Rutilus rutilus). I The effects of constant high temperatures. *Jour. Exp. Bio.* 36(1): 203-216. *Spo. Fish. Abstr.*, 4(2), Abstr. No. 2342, 1959.
- Cocking, A. W., 1959. The effects of high temperature on roach (Rutilus rutilus). II The effects of temperatures increasing to a known constant rate. *Jour. Exp. Bio.*, 36(1): 217-226. *Spo. Fish. Abstr.*, 4(2), Abstr. No. 2344, 1959.
- Coker, R. E., 1933. Influence of temperature on size of fresh water copepods (Cyclops). *Intern. Rev. Ges. Hydrobio. and Hydrograph*, 29(5/6): 406-438.
- Coker, R. E., 1934. Reaction of some fresh water copepods to high temperatures, with a note concerning the rate of development in relation to temperatures. *Jour. Elisha Mitchell Sci. Soc.*, 50(1/2): 143-159.
- Cole, W. H., 1939. The effect of temperature on the color change in Fundulus in response to black and to white backgrounds in fresh and sea water. *Jour. Exp. Zool.*, 80(3): 167-172.
- Collins, G. B., 1952. Factors influencing the orientation of migrating anadromous fishes. *Fish. Bull.*, W.S., 52: 373-396.
- Colton, J. B., 1959. A field observation of mortality of marine fish larvae due to warming. *Limnol. and Oceanogr.*, 4: 219-222.
- Combs, B. D., 1965. Effect of temperature on development of salmon eggs. *Prog. Fish Cult.*, 27(3): 134-137.
- Combs, B. D., and R. E. Burrows, 1957. Threshold temperatures for the normal development of chinook salmon eggs. *Prog. Fish Cult.*, 19(1): 3-6. *Spo. Fish. Abstr.*, 2(3), Abstr. No. 994, 1957.
- Cooper, G. P., and J. L. Fuller, 1945. A biological survey of Moosehead Lake and Hymock Lake, Maine. *Maine Dept. of Inland Fish. and Game, Fish Surv. Rep. No. 6*: 1-160.
- Copeland, J. J., 1936. Yellowstone thermal myxophyceae. *Annals, New York Acad. of Sci.*, 36: 1-232.

- Cory, R. L., and H. F. Davis, 1965. Automatic data system aids thermal pollution study of Patuxent River. *Water and Sewage Works*, 112(4):129.
- Cotter, T. J., and A. W. Lotz, 1961. Cooling pond design in the southwest. *Jour. of the Power Div., ASCE*, 87(P02), Proc. Paper 2861, pp. 85-103.
- Courrier, R., 1922. Seu l'indépendance de la glande seminale et des caracteres sexuels secondaires chez les poissons. *C. R. Acad. Sci. Paris*, 174:70.
- Coutant, C. C., 1962. The effect of a heated water effluent upon the macroinvertebrate riffle fauna of the Delaware River. *Penn. Acad. of Sci.*, 37:58-71.
- Coutant, C. C., 1965. Effect of thermal pollution on Delaware River invertebrates. *Minutes, Third Annual Conf. on the Patuxent Estuary Studies*, Nov. 13-14, 1964. *Chesapeake Bio. Lab. Ref. No.* 65-23.
- Craigie, D. E., 1963. An effect of water hardness in the thermal resistance of the rainbow trout, Salmo gairdnerii Richardson. *Can. Jour. Zool.*, 41(5): 825-830.
- Craig-Bennet, A., 1930. The reproductive cycle of the three-spined stickleback. *Phil. Trans.*, B119:197.
- Crawford, D. R., 1930. Some considerations in the study of the effects of heat and light on fishes. *Copeia*, 73:89-92.
- Creaser, C. W., 1930. Relative importance of hydrogen-ion concentration, temperature, dissolved oxygen and carbon dioxide on habitat selection by brook trout. *Ecol.*, 11:246-262.
- Crisp, D. J., 1957. Effect of low temperature on breeding of marine animals. *Nature, London*, 179:1138-1139.
- Crisp, D. J., (Ed.), 1964. The effects of the severe winter of 1962-63 on marine life in Britain. *Jour. Animal Ecol.*, 33:165-210.
- Cushing, C. E., Jr., and P. A. Olson, 1936. Effects of weed burning on stream conditions. *Trans. Am. Fish. Soc.*, 92(3):303-305.
- Czeczuga, B., 1959. On oxygen minimum and maximum in the metalimnion of Rajgrad Lakes. *Acta Hydrobio.*, 1(2): 109-122.
- Dakin, W. J., 1912. Aquatic animals and their environment. The constitution of the external medium and its effect upon the blood. *Int. Rev. Hydrobio. Hydrog.*, 5:53-80.
- Dallinger, W. H., 1887. Acclimatization of flagellates to high temperatures. *Jour. Roy. Microbiol. Soc., London*, 7:185-199.

- Damann, K. E., 1941. Quantitative study of the phytoplankton of Lake Michigan at Evanston, Illinois. *Butler Univ. Bot. Studies*, 5(1/8): 27-44.
- Dannevig, H., 1894. The influence of temperature on the development of the eggs of fishes. *Rep. Fish. Bd. Scot.*, 13: 147-153.
- Davenport, C. B., and W. E. Castle, 1895. Studies on morphogenesis. III On the acclimatization of organisms to high temperatures. *Arch. Entra. Mech. Org.*, 2: 227-249.
- Davidson, C. M., M. Boothroyd, and D. L. Georgala, 1966. Thermal resistance of Salmonella senftenberg. *Nature*, 212(5066): 1060-1061.
- Davis, R. E., 1955. Heat transfer in the goldfish, Carassius auritus. *Copeia*, 13: 207-209.
- Deelder, C. L., 1958. Some remarks on the stay of fishes under and in ice. *Natura*, 55: 4-6. *Spo. Fish. Abstr.*, 5(3), 1960.
- Deissler, R., 1962. Turbulence in the presence of a vertical body force and temperature gradient. *Jour. Geophysical Resrch.*, 67(8), July.
- Delay, W. H., 1963. Temperature studies on the Umpqua River, Oregon. Water temperature, influences, effects, and control. *Proc. of the 12th Pacific Northwest Symposium on Water Poll. Resrch.*, Pacific Northwest Lab., Publ. Health Serv., U. S. Dept. of Health, Education and Welfare.
- Dendy, J. S., 1944. Further studies of depth distribution of fish, Norris Reservoir, Tennessee. *Jour. Tenn. Acad. Sci.*, 21(1): 94-104.
- Dendy, J. S., 1945. Predicting depth distribution of fish in three TVA storage type reservoirs. *Trans. Am. Fish. Soc.*, 75: 65-71.
- Dendy, J. S., 1946. Water temperature and spring fishing. Norris Reservoir, Tennessee. *Jour. Tenn. Acad. Sci.*, 21: 89-93.
- Dendy, J. S., and R. H. Stroud, 1949. The dominating influence of Fontana Reservoir on temperature and dissolved oxygen in the Little Tennessee River and its impoundments. *Jour. of Tenn. Acad. of Sci.*, 24(1): 41-51.
- Denys, H. A., and J. M. Joseph, 1956. Relationships between temperature and blood oxygen in the large mouth bass. *Jour. Wildl. Mgmt.*, 20(1): 156-164.
- DeRoche, S. E., and L. H. Bond, 1955. Lake trout of Cold Stream Pond, Enfield, Maine. *Trans. Am. Fish. Soc.*, 85: 257-270.

- DeTurville, C. M., and R. T. Jarman, 1965. The mixing of warm water from the Uskmouth Power Stations in the estuary of the River Usk. Int. Jour. of Air and Water Poll., Pergamon Press, England, 9:239-251.
- DeWitt, R. M., 1955. Ecology and life history of the pond snail, Physa gyrina. Ecol., 36(1): 40-44.
- Dickie, L. M., 1959. Water temperature and survival of giant scallop, Trans. Am. Fish. Soc., 88(1): 73.
- Dildine, G. C., 1936. The effect of light and heat on the gonads of Lebistes. Anat. Record 67, Suppl. 1: 61.
- Dimick, R. E., and F. Merryfield, 1945. Fishes of the Willamette River System in relation to pollution. Oregon State College Engng. Exp. Sta. Bull., Ser. 20.
- Donaldson, L. R., and F. J. Foster, 1941. Experimental study of the effect of various water temperatures on the growth, food-utilization, and mortality rate of fingerling sockeye salmon. Trans. Am. Fish. Soc., 70: 339-346.
- Dodd, J. M., and J. N. Dent, 1963. Thyroid gland and temperature tolerance relationships in cold-blooded vertebrates. Nature, London, 199: 299.
- Dorris, T. C., B. J. Copeland, and G. J. Lauer, 1963. Limnology of the middle Mississippi River. IV Physical and chemical limnology of river and chute. Limnol. and Oceano., 8(1): 79-88.
- Doty, M. S., and M. Oguri, 1957. Evidence for a photosynthetic daily periodicity. Limnol. and Oceano., 2(1): 37-40.
- Doudoroff, P., 1938. Reactions of marine fishes to temperature gradients. Bio. Bull., 75: 494-509.
- Doudoroff, P., 1942. The resistance and acclimatization of marine fishes to temperature changes. I Experiments with Girella nigricans (Ayres). Bio. Bull., 83: 219-244.
- Doudoroff, P., 1945. The resistance and acclimatization of marine fishes to temperature changes. II Experiments with Fundulus and Atherinops. Bio. Bull., 88(2): 194-206.
- Downing, K. M., and C. J. Merckens, 1957. The influence of temperatures on the survival of several species of fish in low tensions of dissolved oxygen. Ann. Appl. Bio., 45(2): 261-267.
- Dryer, W. and N. G. Benson, 1957. Observations on the influence of the new Johnsonville steam plant on fish and plankton populations. Tenn. Game and Fish Comm. Rep., 10th Ann. Conf. Southeast Asso. of Game and Fish Comm., pp 85-91. Spo. Fish. Abstr., 2(4), 1957.

Dunnington, E. A., 1965. The effect of heat on attachment of fouling organisms. Minutes of the Third Ann. Conf. on the Patuxent Estuary Studies, Nov. 13-14, 1964. Chesapeake Bio. Lab. Ref. No. 65-23.

Dutton, G. J., and J. P. Montgomery, 1958. Gluconamide synthesis in fish and the influence of temperature. Proc. of Biochem. Soc., 70(4): 178.

Duttweiler, D. W., 1963. A mathematical model of stream temperature. Ph.D. Thesis, The Johns Hopkins Univ., Baltimore, Md., 1963.

Duttweiler, D. W., 1961. Periodic variation of temperature in natural waters. Johns Hopkins Univ. Dept. Sanit. Engng. Water Resour., Low-flow Augmentation Project, Rep. XII: 56-62.

Dysart, B. C., III, and P. A. Krenkel, 1965. The effects of heat on water quality. Proc. of the 20th Indus. Waste Conf., Purdue Univ., Lafayette, Indiana, XLIX(4): 18-39.

Dzyan, Y., 1959. Changes in the temperature preferences of certain species of sturgeon at varying levels of food supply. Nauch. Soobscheniya Inst. Fiziol. Akad. Nauk SSSR 1: 125-127. Referat. Zhur., Bio., 1961, No. 4D74. Spo. Fish Abstr., 7(2), 1962.

Echo, J. B., 1955. Some ecological relationships between yellow perch and cutthroat trout in Thompson Lakes, Montana. Trans. Am. Fish. Soc., 84: 239-248.

Edinger, J. E., and J. C. Geyer, 1965. Heat exchange in the environment. Edison Electric Institute Publ. No. 65-902.

Edmonson, C. H., 1929. Hawaiian Atyidae. Bernice P. Bishop Museum of Polynesian Ethnol. and Nat. Hist. Bull., 66: 3-36.

Edwards, G. A., and L. Irving, 1943. The influence of temperature and season upon oxygen consumption of the sand crab, Emerita talpoida. Jour. Cell. Comp. Physiol., 21: 169-182.

Ekberg, D. R., 1962. Anaerobic and aerobic metabolism in gills of the crucian carp adapted to high and low temperatures. Comp. Biochem. Physiol., 5: 123-128.

Elder, R. A., 1958. Thermal density underflow design and experience. Engng. Lab., Div. of Water Control Planning, Tenn. Valley Author., Norris, Tenn., July.

Elder, R. A., 1959. Thermal density currents and their utilization in condensing water design. Engng. Lab., Div. of Water Control Planning, Tenn. Valley Author., Norris, Tenn., April.

- Elder, R. A., and G. B. Dougherty, 1958. Thermal density underflow diversion, Kingston Steam Plant. Jour. of the Hydraulics Div., ASCE, 84(HY2), Proc. Paper 1583.
- Eldridge, E. F., 1960. The effect of irrigation on the quality of water for industrial use. Proc. 14th Indus. Waste Conf., Purdue Univ., Lafayette, Indiana, p. 52.
- Ellis, M. M., 1942. Fresh-water impoundments. Trans. Am. Fish. Soc., 71(1941): 80-93.
- Ellis, M. M., 1947. Temperature and fishes. U. S. Fish and Wildl. Serv., Fish. Leaflet 221.
- Elser, H. J., 1965. Effect of a warmed-water discharge on angling in the Potomac River, Maryland, 1961-62. Prog. Fish. Cult., 27(2): 79-86. Spo. Fish. Abstr., 10(2), Abstr. No. 7669, 1965. Water Poll. Abstr., 39(2), Abstr. No. 330, 1966.
- Elson, P. F., 1940. Effects of current on the movement of speckled trout. Jour. Fish. Resch. Bd. Can., 4: 517-520.
- Elson, P. F., 1942. Effect of temperature on activity of Salvelinus fontinalis. Jour. Fish. Resch. Bd. Can., 5: 461-470.
- Embody, G. C., 1934. Relation of temperature to the incubation periods of four species of trout. Trans. Am. Fish. Soc., 64: 281-292.
- Engle, M. D., 1960. Condensing water. How does it affect the river? ASME Paper No. 60-WA-92.
- Enropeyzena, N. V., 1944. Preferred temperatures of fish larvae. C. R. Acad. Sci., Moscow, N. S., 42(3): 138-142.
- Evans, R. G., 1948. The lethal temperatures of some common British littoral molluscs. Jour. Animal Ecol., 17: 165-173.
- Evans, R. M., F. C. Purdie, and C. P. Hickman, Jr., 1962. The effect of temperature and photoperiod on the respiratory metabolism of rainbow trout (Salmo gairdneri). Can. Jour. Zool., 40: 107-118.
- Fabricius, E., 1950. Heterogeneous stimulus summation in the release of spawning activities in fish. Inst. Freshwater Res. Drottningholm, 31: 59-99.
- Falkner, N. W., and A. H. Houston, 1966. Some haematological responses to sublethal thermal shock in the goldfish, Carassius auratus L. Jour. Fish. Resch. Bd. Can., 23(8): 1109-1120.

Farmanfarmaian, A., and A. C. Giese, 1963. Thermal tolerance and acclimation in the western purple sea urchin, Strongylocentrotus purpuratus. *Physiol. Zool.*, 36: 237-243.

Farris, D. A., 1961. Abundance and distribution of eggs and larvae and survival of larvae of jack mackerel (Trachurus symmetricus). *Fish. Bull. U. S.*, 61(187): 247-279.

Ferguson, R. G., 1958. The preferred temperature of fish and their midsummer distribution in temperate lakes and streams. *Jour. Fish. Resch. Bd. Can.*, 15: 607-624. *Spo. Fish. Abstr.*, 4(1), Abstr. No. 2275, 1959.

Fish, C. J., 1927. Production and distribution of cod eggs in Massachusetts Bay in 1924 and 1925. *Bull. M. S. Buhr, Fish* 43 (Part III); 253-296.

Fish, F. F., 1948. The return of blueback salmon to the Columbia River. *Sci. Month. (Washington)*, 66: 283-292.

Fisher, C., and C. M. Sullivan, 1958. The effect of temperature on the spontaneous activity of speckled trout before and after various lesions of the brain. *Can. Jour. of Zool.*, 36(1): 49-63.

Fisher, K. C., 1958. An approach to the organ and cellular physiology of adaptation to temperature in fish and small mammals. *Physiol. Adapt.* (Prosser, C. L., ed.), Washington, Am. Physiol. Soc., Inc.

Fisher, K. C., and P. F. Elson, 1950. The selected temperature of Atlantic salmon and speckled trout and the effect of temperature on the response to an electrical stimulus. *Physiol. Zool.*, 23: 27-34.

Fleming, R. H., 1956. The influence of hydrographic conditions on the behavior of fish. *Fish. Bull. F.A.O.* 9: 181-196.

Foerster, R. E., 1937. The relation of temperature to the seaward migration of young sockeye salmon and speckled trout and the effect of temperature on the response to an electrical stimulus. *Physiol. Zool.*, 23: 27-34.

Fortune, P. Y., 1958. The effect of temperature changes on the thyroid-pituitary relationship in teleosts. *Jour. Exp. Bio.*, 35(4): 824-831.

Fox, D. L., and E. F. Corcoran, 1957. Thermal and osmotic countermeasures against some typical marine fouling organisms. *Corrosion*, 14: 31-32.

Fox, H. M., 1936. The activity and metabolism of poikilothermal animals in different latitudes, I. *Proc. Zool. Soc. London*, 945-955.

Fox, H. M., 1938. The activity and metabolism of poikilothermal animals in different latitudes, III. *Proc. Zool. Soc. London*, 11C8, 501-505.

- Fox, H. M., 1939. The activity and metabolism of poikilothermal animals in different latitudes, V. Proc. Zool. Soc. London A., 109, 141-156.
- Fox, H. M., and C. A. Wingfield, 1937. The activity and metabolism of poikilothermal animals in different latitudes, II. Proc. Zool. Soc. London A., 107, 275-282.
- Fraenkel, G., and D. L. Gunn, 1940. The orientation of animals: Temperature reactions. Oxford Clarendon, 352 pp, Chap. 14.
- Fraenkel, G., and H. S. Hapf, 1940. The physiological action of abnormally high temperatures in poikilothermic animals. I Temperature adaptation and the degree of saturation of the phosphatides. Biochem. Jour., 34(7): 1085-1092.
- Freeman, J. A., 1950. Oxygen consumption, brain metabolism, and respiratory movements of goldfish during temperature acclimatization, with special reference to lowered temperatures. Bio. Bull., 99: 416-424.
- Fries, E. F. B., 1952. Observations on chill tolerance and applied chill coma in subtropical American marine fishes, especially. Copeia, (3): 147-152.
- Fry, A. S., M. A. Churchill, and R. A. Elder, 1963. Significant effects of density currents in TVA's integrated reservoir and river system. Proc., Minnesota International Hydraulics Conv., Minneapolis, Minnesota, Sept.
- Fry, F. E. J., 1947. Effects of the environment on animal activity. Univ. of Toronto Study Bio. Ser. 55: Publ. Ont. Fish. Resch. Lab., 68: 1-62.
- Fry, F. E. J., 1951. Some environmental relations of the speckled trout (Salvelinus fontinalis). Proc. Northeast Atlantic Fish. Conf., May 1951: 1-29.
- Fry, F. E. J., 1951. Some temperature relations of fish. Fed. Proc., 10(1): 46.
- Fry, F. E. J., 1957. The lethal temperature as a tool in taxonomy. Anee Biologique, 33(5-6): 205-219.
- Fry, F. E. J., 1958. Temperature compensation. Ann. Rev. Physiol., 20: 207-224.
- Fry, F. E. J., 1964. Animals in aquatic environments: fishes. Handbook of Physiol. (Dill, D. B., ed.), Sec. 4, Adaptation to the Environment: 715-728. Am. Physiol. Soc., Wash., D.C.

- Fry, F. E. J., V. S. Black, and E. C. Black, 1947. Influence of temperature on the asphyxiation of young goldfish (Carassius auratus L.) under various tensions of oxygen and CO₂. Bio. Bull., 92: 217-224.
- Fry, F. E. J., J. R. Brett, and G. H. Clawson, 1942. Lethal limits of temperature for young goldfish. Rev. Can. Bio., 1: 50-56.
- Fry, F. E. J., et al, 1942. Temperature acclimatization. Fish. Rev. Can. Bio., 1: 50-56.
- Fry, F. E. J., and O. Fournier, 1942. Les températures lethales de divers organismes aquatiques du haut Saint - Laurent. Res. Can. Bio., 1: 103-104.
- Fry, F. E. J., and J. S. Hart, 1946. The relation of temperature to oxygen consumption in the goldfish. Bio. Bull., 94(1): 66-77.
- Fry, F. E. J., and J. S. Hart, 1948. Cruising speed of goldfish in relation to water temperature. Jour. Fish. Resch. Bd. Can., 7: 169-175.
- Fry, F. E. J., J. S. Hart, and K. F. Walker, 1946. Lethal temperature relations for a sample of young speckled trout, Salvelinus fontinalis. Univ. Toronto Stud. Bio. Ser. 54, Publ. Ont. Fish. Resch. Lab., 66: 1-35.
- Fuhrman, F. A., N. Hollinger, J. M. Crimson, J. Field, and F. W. Weymouth, 1944. The metabolism of the excised brain of the large-mouthed bass, (at graded temperature levels). Physiol. Zool., 17: 42-50.
- Galloway, J. C., 1951. Lethal effects of the cold winter of 1939/40 on marine fishes at Key West, Florida. Copeia, 2: 118-119.
- Gameson, A. L. H., J. W. Gibbs, and M. J. Barrett, 1959. A preliminary temperature survey of a heated river. Water and Water Engng., London, 63(755): 13-17.
- Gameson, A. L. H., H. Hall, and W. S. Preddy, 1957. Effects of heated discharges on the temperature of the Thames estuary. Engineer, London 204, 816-9, 850-2, 893-6.
- Gameson, A. L. H., et al, 1961. Effects of heated discharges on the temperature of the Thames estuary - II. Combustion, 32(7): 37.
- Gameson, A. L. H., K. G. Vandyke, and C. G. Ogden, 1958. Effect of temperature on aeration at Weirs. Water and Water Engng., London, 62: 489.
- Garay, P. N., 1966. Design considerations in selecting water cooling systems. Consulting Engineer, 26(1): 92.
- Gardner, J. A., 1926. Report on the respiratory exchange in freshwater fish, with suggestions as to further investigations. Fish. Invest., London, 1(3): 1-17.

- Gardner, J. A., G. King, and E. B. Powers, 1922. The respiratory exchange in freshwater fish III Goldfish. *Biochem. Jour.*, 16: 323-529.
- Garside, E. T., 1959. Some effects of oxygen in relation to temperature on the development of lake trout embryos. *Can. Jour. Zool.*, 37: 689-698.
- Garside, E. T., and J. S. Tait, 1958. Preferred temperature of rainbow trout (Salmo gairdneri Richardson) and its unusual relationship to acclimation temperature. *Can. Jour. Zool.*, 36: 563. *Spo. Fish. Abstr.*, 3(3), Abstr. No. 1676, 1958.
- Gauvin, A. R., 1955. Environmental changes in a polluted stream during winter. *The Am. Midland Naturalist*, 54(1): 78-88.
- Gerbilisky, N. L., 1937. Development of oocytes in Carassius carassius and its dependence upon the temperature. *Bull. Bio. Med. Exp. Moscow* 3: 160-161.
- Gibson, E. S., and F. E. J. Fry, 1954. The performance of the lake trout, Salvelinus namaycush, at various levels of temperature and oxygen pressure. *Can. Jour. of Zool.*, 32: 252-260.
- Gibson, M. B., 1953. Upper lethal temperature relations of the guppy, Lebistes reticulatus. *Can. Jour. Resch. Zool.*, 32: 302-407. *Spo. Fish. Abstr.*, 1(1), Abstr. No. 159, 1955.
- Gibson, M. B., and B. Hirst, 1955. Effect of salinity and temperature on the pre-adult growth of guppies. *Copeia*, 3: 241-243.
- Gieysztor, M., 1960. On the thermal conditions of the littoral zone of lakes. *Polskie Archiwum Hydrobiologii*, VII(XX): 171-193.
- Gieysztor, M., 1961. Temperatures and chemistry of the littoral zone in lakes. *Verh. Int. Ver. Limnol.* 1959, 14: 84-86. *Spo. Fish. Abstr.*, 8(1), 1963.
- Glaser, O., 1929. Temperature and heart rate in Fundulus embryos. *Brit. Jour. Exper. Bio.*, 6: 325-339.
- Goitein, E. E., 1957. Selection and application of cooling towers in steam-electric stations. *Combustion*, 29(5): 38.
- Gotaas, H. B., 1948. Effect of temperature on biochemical oxidation of sewage. *Sewage Works Jour.*, 20(3): 441.
- Gradzinski, Z., 1949. The influence of temperature upon the rate of the heart in the embryos of teleost fishes. *Bull. Int. Acad. Cracovie*, 1948, B.II., 7-10, 255-293.

- Gradzinski, Z., 1950. The susceptibility of the heart in the sea-trout embryo Salmo trutta L. to small changes in temperature. Bull. Intern. Acad. Polonaise. Sci. Ser. BII, 4-6: 173-182.
- Graham, J. M., 1949. Some effects of temperature and oxygen pressure on the metabolism and activity of the speckled trout, Salvelinus fontinalis. Can. Jour. Resrch., 27: 270-288.
- Grainger, J. N. R., 1958. First stages in the adaptation of poikilotherms to temperature change. Physiol. Adapt. Am. Physiol. Soc. (Prosser, E. T., Ed.), Wash., D. C., pp. 79-91.
- Gray, J., 1928. The growth of fish. III The effect of temperature on the development of the eggs of Salmo fario. Brit. Jour. Exp. Bio., 6: 110-130.
- Green, G. H., and F. J. Andrew, 1962. Limnological changes in Seton Lake resulting from hydroelectric diversions. Bio. Abstr., 37, Abstr. No. 2236.
- Greene, C. W., 1930. The smelts of Lake Champlain. Biological Survey of the Champlain Watershed, Suppl. 19th Ann. Rep. New York Conserv. Dept.: 105-129.
- Gunn, D. L., 1942. Body temperature in poikilotherm animals. Bio. Rev. 17: 293-314.
- Gunter, G., 1941. Death of fishes due to cold on the Texas coast, January 1940. Ecol., 22: 203-208.
- Gunter, G., 1957. Temperature. Treatise on Marine Ecology and Palaeoecology I. (Hedgepeth, J. W., Ed.), Chap. 8, Geol. Soc. Am. Mem. 67, 159-184.
- Gutsell, J. S., 1929. Influence of certain water conditions, especially dissolved gases, on trout. Ecol., 10(1): 77-96.
- Halecrow, K., 1963. Acclimation to temperature in the marine copepod, Calanus finmarchicus (Gunner). Limnol. and Oceanogr., 8: 1-8.
- Halsband, E., 1953. Studies of the behavior of trout (Trutta cridea W. Gibb) and chub (Squalius cephalus Heck.) under the influence of various external factors. Zeits. Fisherei, 2:227. Sewage and Indus. Wastes, 27(6), 1955.
- Hancock, H. M., 1954. Investigations and experimentation relative to winter aggregations of fishes in Canton Reservoir, Oklahoma. Resrch. Foundation Publ. No. 58, Okla. Agr. and Mechanical College, Stillwater, Okla.

Harbeck, G. E., Jr., 1953. The use of reservoirs and lakes for the dissipation of heat. Circular No. 282, Geol. Survey, U. S. Dept. of the Interior.

Harbeck, G. E., Jr., et al, 1959. The effect of the addition of heat from a power plant on the thermal structure and evaporation of Lake Colorado City, Texas. Professional Paper No. 272-B, Geol. Survey, U. S. Dept. of the Interior, U. S. Government Printing Office, Washington, D. C.

Harrington, R. W., 1956. An experiment of the effects of contrasting daily photoperiods on gametogenesis and reproduction in the centrarchid fish, Enneacanthus obesus (Girard). Jour. of Exp. Zool., 131: 203-224.

Hart, J. S., 1947. Lethal temperature relations of certain fish of the Toronto region. Trans. Roy. Soc. Can., 41: 57-71.

Hart, J. S., 1952. Geographic variations of some physiological and morphological characters in certain fresh water fish. Univ. Toronto Stud. Bio. Sci. 60, Publ. Ont. Fish Resch. Lab., 72: 1-79.

Hasan, R., and S. Z. Qasim, 1961. Studies on fish metabolism. I, The effect of temperature on the heart rate. Proc. Indiana Acad. Sci., 53B: 230-239.

Haskell, D. C., L. E. Wolf, and L. Bouchard, 1956. The effect of temperature on the growth of brook trout. New York Fish and Game Jour., 3(1): 108-113. Spo. Fish. Abstr., 2(1), 1956.

Haslam, R. I., R. L. Hershey, and R. H. Keen. Effect of gas velocity and temperature on rate absorption. Indus. and Engng. Chem., 16(1924): 1224.

Hata, K., 1950. The influence of environmental changes on freshwater fishes. Studies on the influence of rise of water temperature on young trouts. Bull. Jap. Soc. Sci. Fish., 15(11): 665. Water Poll. Abstr., 26(5): 139, 1953.

Hatfield, H. F., 1960. The effects of the discharge of warm water to rivers. Proc. 21st Ann. Water Conf., Pittsburgh, Pa., p. 102.

Hatfield, H. F., 1962. Temperature distribution in the Susquehanna River at Brunner Island, steam electric station. Prog. Rep. EM-DEV-242.

- Hathaway, E. S., 1927. The relations of temperature to the quantity of food consumed by fish. *Ecol.*, 8: 428-434.
- Hathaway, E. S., 1928. Quantitative study of the changes produced by acclimatization in the tolerance of high temperature by fishes and amphibians. *Bull. U. S. Bur. of Fish.*, 43(2): 169-192.
- Havgaard, N., and L. Irving, 1943. The influence of temperature upon oxygen consumption of the cummer (Tautogolabrus adspersus, Walbaum) in summer and winter. *Jour. Cell. and Comp. Physiol.*, 21: 19-26.
- Hawkes, H. A., 1962. Biological aspects of river pollution. Klein, L. River Poll. II, Causes and Effects. Butterworths, London, pp. 311-432.
- Hayes, F. R., 1949. The growth, general chemistry, and temperature relations of salmonid eggs. *Quart. Rev. Bio.*, 24: 281-308.
- Hayes, F. R., and D. Pelluet, 1945. The effect of temperature on the growth and efficiency of yolk conversion in the salmon embryo. *Can. Jour. Resch.*, Ottawa, 23D: 7-15.
- Hazard, A. S., 1933. Low water temperature, a limiting factor in the successful production of trout in natural waters: *Trans. Am. Fish. Soc.*, 63: 204-207.
- Heath, W. G., 1963. Maximum temperature tolerance as a function of constant temperature acclimation in the Gila topminnow (Poeciliopsis occidentalis). *Diss. Abstr.*, 23: 3559.
- Heath, W. G., 1963. Thermoperiodism in sea-run cutthroat trout (Salmo clarki clarki). *Science*, 142: 486-488.
- Heinicke, E. A., and A. H. Houston, 1965. Effect of thermal acclimation and sublethal heat shock upon ionic regulation in the goldfish Carassius auratus L. *Jour. Fish Resch. Bd. Can.*, 22(6): 1455-1476.
- Hela, I., and T. Laevastu, 1963. The influence of temperature on the behavior of fish. *Arch. Soc. Zool. bot. fenn. Vanamo* 15: 83-103. *Bio. Abstr.*, 42(5), *Abstr. No.* 17941, 1963. *Spo. Fish. Abstr.*, 8(3), *Abstr. No.* 6032, 1963.
- Henderson, C., and R. G. Foster, 1956. Studies of smallmouth black bass (Micropterus dolomieu) in the Columbia River New Richland, Washington. *Trans. Am. Fish. Soc.*, 86: 113-127.
- Henderson, J. T., 1929. Lethal temperatures of Lamellibranchiata. *Contr. Can. Bio. Fish. N. S.* 4: 397-411.
- Henderson, H. E., 1963. Influence of light and temperature on the reproductive cycle of the eastern brook trout, Salvelinus fontinalis (Mitchell). *Jour. Fish. Resch. Bd. Can.*, 20(4): 859-897.

Herry, S., 1959. Pollution of rivers by heated discharges. Bull. Centre. belge Et. Document. Eaux, No. 46: 226-235. Water Poll. Abstr., 36(4), Abstr. No. 686, 1963. Spo. Fish. Abstr., 9(2), Abstr. No. 6589, 1964.

Hewitt, E. R., 1938. Aeration of waters for trout. Prog. Fish Cult., 35: 23.

Higurashi, T., 1925. Optimum water temperature for hatching the eggs of Plecoglossus altivelis. T. and S. J. Imp. Fish. Inst., Tokyo, 20: 12-14.

Higurashi, T., 1925. Optimum water temperature for hatching the eggs of Hypomesus alidus, Pallas. J. Imp. Fish. Inst., Tokyo, 21: 2-5.

Higurashi, T., and M. Tauti, 1925. On the relation between temperature and the rate of development of fish eggs. J. Imp. Fish., Inst., Tokyo, 21: 5-9.

Hildebrand, S. F., and I. Towers, 1927. Food of trout in Fish Lake, Utah. Ecol., 8(4): 389-397.

Hile, R., 1941. Age and growth of the rock bass, Ambloplites rupestris (Rafinesque) in Nebish Lake, Wisconsin. Trans. Wis. Acad. Sci., Arts and Letters, 33: 189-337.

Hile, R., and C. Juday, 1941. Bathymetric distribution of fish in lakes of the northeastern highlands, Wisconsin. Trans. Wis. Acad. Sci., Arts and Letters, 33: 147-187.

Hisaoka, K. K., and C. F. Firlit, 1962. Ovarian cycle and egg production in the zebrafish, Brachydanio rerio. Copeia, 1962: 788-792.

Ho, H. J., 1936. The growth of the goldfish (Carassius auratus). China J. Shanghai, 24: 101-105.

Hoaglund, H., 1932. Impulses from the sensory nerves of catfish. Proc. Nat. Acad. Sci., 18: 700-705.

Hoak, R. D., 1961. Defining thermal pollution. Power Engng., 65: 39-42.

Hoak, R. D., 1961. The thermal pollution problem. Jour. Water Poll. Control Federation, 33(12): 1267-1276.

Hoak, R. D., 1962. The thermal pollution problem. Proc. of the 25th Ann. Short Course, Louisiana St. Univ., Engn. Exp. Sta., Bull. No. 68: 141-146, March 15, 16, and 17, 1962)

- Hoak, R. D., and H. C. Bramer, 1961. Diurnal variation of dissolved oxygen in natural streams. Proc. 16th Ind. Conf., Purdue Univ., pp. 167-175.
- Hoar, W. S., 1942. Diurnal variations in feeding activity of young salmon and trout. Jour. Fish. Resch. Bd. Can., 6(1): 90-101.
- Hoar, W. S., 1955. Seasonal variations in the resistance of goldfish to temperature stress. Trans. Roy. Soc. Can., Ser. 3, Sec. V, No. 49: 25-34.
- Hoar, W. S., 1956. Photoperiodism and thermal resistance of goldfish. Nature, London, 178: 364-365. Spo. Fish. Abstr., 2(4), Abstr. No. 1293, 1957.
- Hoar, W. S., and M. K. Cottle, 1952. Dietary fat and temperature tolerance of goldfish. Can. Jour. Resch. Zool., 30: 41-48.
- Hoar, W. S., and M. K. Cottle, 1952. Some effects of temperature acclimatization on the heat tolerances of goldfish (Carrasius auratus). Can. Jour. Resch., D 27: 85-91.
- Hoar, W. S., and J. E. C. Dorchester, 1949. Effect of dietary fat on the heat tolerance of goldfish (Carrasius auratus). Can. Jour. Resch., 27: 85-91.
- Hoar, W. S., and G. B. Robertson, 1959. Temperature resistance of goldfish maintained under controlled photoperiods. Can. Jour. Zool., 37: 419-428. Spo. Fish. Abstr., 4(4), Abstr. No. 2743, 1959.
- Hobbie, J. E., 1961. Summer temperatures in Lake Schrader, Alaska. Limnol. and Oceano., 6(3): 326-329.
- Hochachka, P. W., and F. R. Hayes, 1962. The effect of temperature acclimation on pathways of glucose metabolism in the trout. Can. Jour. Zool., 40(2): 261-270.
- Hockley, A. R., 1963. Some effects of warm water effluents in Southampton Water. Ann. Rep. Challenger Soc., 3(15): 37-38.
- Hoestlant, H., and J. Lasblier, 1959. Thermal destruction of the fresh water mussel. Destruction thermique de la moule d'eau douce. L'Eau, 46: 259, 1959. Publ. Health Engng. Abstr., 40(7): 35, 1960. Water Poll. Abstr., 34(12): 404, 1961.
- Holton, G. D., 1953. A trout population study on a small creek in Gallath County, Montana. Jour. Wildl. Mgmt., 117: 62-82.
- Hornyold, A. G., 1926. Reduction of the body size of eels during development of pigmentation. Notos y Resumenes, Inst. Espanol. Oceano., 10: 1-8.

Houston, A. H., 1962. Some observations on water balance in the goldfish, Carassius auratus L., during cold death. Can. Jour. Zool., 40(7): 1169-1174.

Hubbs, C. L., 1923. Seasonal variation in the number of vertebrae of fishes. Papers Mich. Acad. Sci., 2: 217-224.

Hubbs, C., 1951. Minimum temperature tolerances for fishes of the genera Signalosa and Herichthys in Texas. Copeia, 4: 297.

Hubbs, C., 1961. Developmental temperature tolerances of four etheostomatine fishes occurring in Texas. Copeia, 1961: 195-198.

Hubbs, C., 1964. Effects of thermal fluctuations on the relative survival of greenthroat darter young from stenothermal and eurythermal waters. Ecol., 45(2): 376-379.

Hubbs, C., 1965. Developmental temperature tolerance and rates of four southern California fishes, Fundulus parvipinnis, Atherinops affinis, Leuresthes tenuis, and Hypsoblennius sp. Calif. Fish and Game, 51(2): 113-122.

Hubbs, C., and N. E. Armstrong, 1962. Developmental temperature tolerance of Texas and Arkansas-Missouri Etheostoma spectabile (Percidae, Osteichthyes). Ecol., 43: 742-744.

Hubbs, C., and W. F. Hettler, 1964. Observations on the toleration of high temperatures and low dissolved oxygen in natural waters by Crenichthys baileyi. S. West Nat., 9: 245-248. Bio. Abstr. 46, Abstr. No. 3744, 1965. Water Poll. Abstr., 39(1), Abstr. No. 34, 1966.

Hubbs, C., and K. Strawn, 1957. The effects of light and temperature on the fecundity of the greenthroat darter, Etheostoma lepidum. Ecol., 38(4): 596. Spo. Fish. Abstr., 3(1), Abstr. No. 1442, 1958.

Huet, M., 1965. Water quality criteria for fish life. pp. 160-167. Biological Problems in Water Poll., Third Seminar, 1962 (Tarzwell, C. M., Ed.), Robert A. Taft Engng. Ctr., Publ. Health Serv., Publ. No. 999-WP-25, 434 pp.

Huntsman, A. G., 1942. Death of salmon and trout with high temperature. Jour. Fish. Resch. Bd. Can., 5: 485-501.

Huntsman, A. G., 1946. Heat stroke in Canadian maritime stream fishes. Jour. Fish Resch. Bd. Can., 6: 476-482.

Huntsman, A. G., and M. I. Sparks, 1924. Limiting factors for marine animals 3, Relative resistance to high temperature. Contr. Can. Bio., 2: 102-113.

Hutchinson, L., 1947. Analysis of the activity of the fresh water snail, Viviparus malleatus (Reeve). Ecol., 28(4): 335-345.

Hynes, H. B. N., 1959. The biological effects of water pollution. The Effects of Poll. on Living Material (Yapp, W. B., Ed.), London, p. 11.

Hynes, H. B. N., 1963. The biology of polluted waters. Liverpool Univ. Press, 202 pp., Chap. 9, Heat, Salts, and Pollution of Lakes: 136-145.

Ide, F. P., 1935. The effect of temperature on the distribution of the mayfly fauna of a stream. Univ. Tor. Stud. Bio. Ser., 39: 9-76.

Iezzi, T., J. A. Filson, and C. S. Myers, 1952. Effects of temperature changes and temperature levels on Lepomis macrochirus macrochirus (bluegill sunfish) and on Salmo fario (brown trout). Penn. Dept. Publ. Health, Indus. Wastes Div. (mimeo rep.).

Iles, R. B., 1963. Cultivating fish for food and sport in power station water. New Scientist, 117: 227-229.

Ingleson, H., and A. Harrison, 1941. Effect of temperature on the exchange capacities of some base-exchange materials used in water softening. Jour., Soc. of Chem. Indus., 60: 87. Water Poll. Abstr., 14, June.

Ingram, W. M., and W. W. Towne, 1959. Stream life below industrial outfalls. Publ. Health Rep., 74: 12.

Irving, L., E. C. Black, and V. Safford, 1941. The influence of temperature upon the combination of oxygen with the blood of trout. Bio. Bull., 80: 1-17.

Irvine, D. G., K. Newman, and W. S. Hoar, 1957. Effects of dietary phospholipid and cholesterol on the temperature resistance of goldfish. Can. Jour. Zool., 35(6): 691-709. Spo. Fish. Abstr., 3(2), Abstr. No. 1585, 1958.

Ivanova, L. V., 1962. The role of thermokinesis in the distribution of animals within the temperature gradient (based on an analysis of the temperature reactions of Anopheles plumbeus larvae). Zool. Zhur., 41(9): 1345-1352. Bio. Abstr., 42(4), No. 14147, 1963.

Ivlev, V. S., 1958. Ecological and physiological analysis of fish distribution in environmental gradients. Tr. Soveshch. Ikhtiolog. Komis. Akad. Nauk SSSR 8: 288-296. Referat. Zhur. Bio. 1959, No. 71778. Bio. Abstr., 44(3), Abstr. No. 9265, 1963. Spo. Fish. Abstr., 9(1), Abstr. No. 6425, 1964.

James, M. C., O. L. Meehan, and E. J. Douglass, 1944. Fish stocking as related to the management of inland waters. U. S. Fish Wildl. Serv. Conserv. Bull. 35, 22 pp.

Jammes, L., 1931. Sur le compartiment, au lac d'aredon, de la truite commune Salmo trutta L. en fonction des agents thermique et nutritifs. C. R. Soc. Bio., Paris, 107: 1482-1485.

John, K. R., 1956. Onset of spawning activities of the shallow water cisco, Leucichthys artedi (LeSuer), in Lake Mendota, Wisconsin, relative to water temperatures. Copeia, 1956 (2): 116-118.

Johnson, F. H., 1957. Influence of temperature on biological systems. Am. Physiol. Soc., Washington, D. C., 275 pp. (ed.).

Johnson, M. G., and W. H. Charlton, 1960. Some effect of temperature on the metabolism and activity of the largemouth bass, Micropterus salmoides Lacepede. Prog. Fish. Cult., 22: 155. Spo. Fish. Abstr., 6(1), Abstr. No. 3890, 1961.

Jones, J. R. E., 1949. An ecological study of the river Rheidol, North Cardiganshire, Wales. Jour. Animal Ecol., 18: 67-88.

Jones, J. R. E., 1962. Fish and river pollution. River Pollution, II. Causes and Effects (Klein, L., Ed.): 255-310, Butterworth, London.

Jones, J. R. E., 1964. Thermal pollution: the effect of heated effluents. Fish and River Pollution, Chap. 13: 153-168, Butterworth and Co., Ltd., Washington, D.C.

Kawajiri, M., 1927. The optimum temperature of water for the hatching of eggs of trout, Oncorhynchus masacu (Walbaum). Jour. Imp. Fish. Inst., 3: 14-18.

Kawajiri, M., 1927. On the optimum temperature of water for hatching the eggs of rainbow trout (Salmo irideus, Gibbon). Jour. Imp. Fish. Inst., 23: 59-64.

Kawajiri, M., 1928. The influence of variation of temperature of water on the development of fish eggs. On the relation of growth and death from starvation of the trout fry to temperature. On the studies of the population - density of cultured fishes. Jour. Imp. Fish. Inst., Tokyo, 24: 1-12.

Kazman, R. G., 1948. River infiltration as a source of ground water supply: (a) temperature characteristics of infiltrated water; (b) Fig., 6, thermographs of Ohio River water and pumped water entering a reservoir. Transactions, ASCE, 113.

- Keiz, G. Uder den temperatureinfluss auf die erbrutung der eier des karpfens (Cyprinus carpio L., v. Vertebrata Pisces). Naturwiss, 46: 499.
- Kennedy, W. A., 1940. The migration of fish from a shallow to a deep lake in spring and early summer. Trans. Am. Fish. Soc., 70: 391-396.
- Kerr, J. E., 1953. Studies on fish preservation at the Contra Costa Steam Plant of the Pacific Gas and Electric Company. Fish. Bull. No. 92, Calif. Dept. of Fish and Game.
- Kerr, S. L., and S. Moyer, 1962. Hydraulic-engineering problems at Southwork Generating Station. Transactions, Am. Soc. of Mech. Engrs., 64(6): 529, Aug.
- Ketchen, K. S., 1952. Factors influencing the survival of the lemon sole (Parophrys netulus, Girard) in Hecati Strait, British Columbia. Univ. Toronto, Dept. of Zool., Ph.D. Thesis.
- Kinne, O., 1956. Über temperatur und salzgehalt and ihre physiologisch-biologische bedeutung. Bio. Zbl. 75: 314-327.
- Kinne, O., 1956. Über den einfluss des salzgehaltes und der temperatur auf wachstum, form und vermehrung bei den hydroidopolypen Cordylophora caspia (Pallas), thecata, claudidae. Zool. Jb. (Physiol.) 66: 565-638.
- Kinne, O., 1960. Growth, food uptake, and food conversion in a euryplastic fish exposed to different temperatures and salinities. Physiol Zool. 33: 288-317.
- Kinne, O., 1963. The effects of temperature and salinity on marine and brackish water animals. I. Temperature. Oceano. Mar. Bio. Ann. Rev. 1: 301-340.
- Kinne, O., 1964. The effects of temperature and salinity on marine and brackish water animals. II. Salinity and temperature salinity combinations. Oceano. Mar. Bio. Ann. Rev. 2: 281-339.
- Kinne, O., 1965. Salinity requirements of the fish, Cyprinodon mascularius. Biological Problems in Water Pollution, Transactions of the 3rd Seminar (1962), Robert A. Taft Engng. Center, Cincinnati, Ohio, U. S. Publ. Health Serv., Publ, 999-WP-25: 187-192.
- Kinne, O., and E. M. Kinne, 1962. Rates of development in embryos of a cyprinodont fish exposed to different temperature - salinity - oxygen combinations. Can. Jour. Zool., 40(2): 231-253.
- King, J. E., 1943. Survival time of trout in relation to occurrence. Am. Md. Nat., 29: 624-642.

- Klein, L., 1962. River pollution II. Causes and effects. Butterworth, London, 456 pp.
- Knight, A. W., 1963. The effect of water flow, temperature and oxygen concentration on the plecoptera nymph, Acroneuria Pacifica Banks. Proc. Utah Acad. Sci., Arts and Letters, 40, Part II.
- Knight, A. W., 1964. Relative importance of varying oxygen concentration, temperature, and water flow on the mechanical activity and survival of the plecoptera nymph, Pteronarcys Califronica Newport. Proc. Utah Acad. Sci., Arts and Letters, 41, Part I.
- Kowalska, A., 1959. On the influence of temperature on the embryonic development of the brook trout (Salmo trutta m. fario L.). Przegl. Zool., 3:253-259.
- Kropp, B. N., 1947. The effect of temperature on the rate and variation of opercular movement in Fundulus diaphanous diaphanous. Can. Jour. Resch., 25D, 2:91-95.
- Krough, A., 1939. Osmoregulation in aquatic animals. Univ. Press Cambridge, 242 pp.
- Kruppert, H. H., and P. D. Meijering, 1963. Die beziehungen zwischen temperatur und hell-dunkel-farbwechsel bei der scholle (Pleuronectes platessa L.). Zool. Anz. 170:55-61.
- Kubo, T., 1936. Feeding velocity of the eel, Anguilla japonica (TandS) in relation to water temperature and other environmental conditions. Bull. Jap. Soc. Sci. Fish., 4:335-338.
- Kubo, T., 1953. On the blood of salmonid fishes of Japan during migration. I. Freezing point of blood. Bull. Fac. Fish. Hakhaida Univ., 4:138-148.
- Kujiyama, E., 1929. On the influence of temperature upon the development of eggs of Pograsomas major (TandS). Jour. Imp. Fish. Inst. Tokyo, 24:109-113.
- Kuroki, T., 1954. On the relation between water temperature and the response for stimuli. The investigation to decide the "optimum temperature." Mem. Fac. Fish., Kagoshima Univ., 3(2):19-24. Spo. Fish. Abstr., 4(3), 1959.
- Kuthalingam, M.D.K., 1959. Temperature tolerance of the larva of ten species of marine fishes. Curr. Sci. 28:75-76.
- Leberge, R. H., 1959. A critical problem in stream pollution comes from thermal discharges. Water and Sewage Works, 106:536-540.

- Lagerspetz, K., and V. Perttunen, 1962. Effect of temperature on the periodic heart beat reversal and heart rate in Corethra plumicornis (Diptera). Jour. Insect Physiol., 8(11-12): 621-625. Bio. Abstr., 42(2), No. 8149, 1963.
- Langbein, W. B., et al, 1949. Annual runoff in the United States. U. S. Geol. Surv. Circ. 52.
- Langhear, J. W., 1953. Cooling ponds may answer your water cooling problem. Chem. Engng., pp. 194-198, Aug.
- Larkin, P. A., 1958. The investigation of fish-power problems. Univ. of Brit. Columbia, Vancouver, 111 pp.
- Lawrence, W. M., 1940. The effect of temperature on the weight of fasting rainbow trout fingerlings. Trans. Am. Fish. Soc., 70: 290-296.
- LeBosquet, M., Jr., 1946. Cooling water benefits from increased river flows. Jour. New England Water Works Asso., 60(2): 111-116.
- LeBourueau, J. W., 1959. Second report of Deerfield River study by Yankee Atomic Electric Co., Yankee Atomic Electric Co., 441 Stuart St., Boston, Mass.
- Lewis, A. G., 1959. The vertical distribution of some inshore copepods in relation to experimentally produced conditions, of light and temperature. Bull. Mar. Sci. Gulf and Caribbean, 9(1): 69-78.
- Lewis, R. M., 1965. The effect of minimum temperature on the survival of larval Atlantic menhaden, Brevoortia tyrannus. Trans. Am. Fish. Soc., 94: 409-412.
- Liebmann, H., 1958-60. Handbook of freshwater and waste water biology. 2(1-5): 960. Bio. Abstr. 36(2), Abstr. No. 3394, 1961. Spo. Fish. Abstr., 6(2), Abstr. No. 4179, 1961.
- Ljubtzky, A. I., 1935. Zur erforschung des temperatureffects in du morphogenese. II. Einfluss der temperatur auf die entwiclungs-geschwindigkeit und wachstrum des embryos von Salmo trutta, fario. Zool. Jahr B. Jena (Allg. Zool.), 54: 405-422.
- Lloyd, R., and D. W. M. Herbert, 1962. The effect of the environment on the toxicity of poisons to fish. Jour. Institute of Publ. Health Engnrs., pp. 132-145.
- Loeb, H. A., 1960. Survival of carp at a low temperature. New York Fish and Game Jour., 7(2): 158-159.

- Loeb, H. A., 1964. Submergence of brown bullheads in bottom sediments. New York Fish and Game Jour., II(2): 119-124.
- Loeb, J., and H. Wastenys, 1912. On the adaptation of fish (Fundulus) to high temperature. Jour. of Exp. Zool., 12: 543-557.
- Long, L. K., N. D. Zau, and N. K. Ving, 1961. First results of a study of the action of temperature on the fish, Tilapia mossambica Peters, acclimated in Vietnam since 1961. Zhur Coshei Bio., 22(6): 444-451. Referat. Zhur., Bio. 1962, No. 9163 (Transl.). Bio. Abstr., 41(3), 1962.
- Luferova (Lokhanina), L. A., 1960. The influence of hydro-electric power station on the zooplankton in Gorky reservoir. Byulletin Instituta Biologii Vodokhranilischeh, Akad. Nauk, USSR, 6: 38-39.
- Lund, J. W. G., and J. F. Talling, 1957. Botanical limnological methods with special reference to the algae. The Botan. Rev., 23: 489-583.
- Macan, T. T., 1959. The temperature of a small stony stream. Hydrobiologia, Freshwater Bio. Asso., Ambleside, England, Vol. 12.
- MacArthur, J. W., and W. N. T. Baillie, 1929. Metabolic activity and duration of life. I-Influence of temperature on longevity in Daphnia magna. Jour. Exp. Zool., 53(2): 221-242.
- MacCardle, R. C., 1937. The effect of temperature on mitochondria in liner cells of fish. Jour. Morphol., 61: 613-639.
- Maciolek, J. A., and P. R. Needham, 1951. Ecological effects of winter conditions on trout and trout foods in Convict Creek, California. Trans. Am. Fish. Soc., 81: 202-217.
- MacKichan, K. A., 1957. Estimated use of water in the United States, 1955. U. S. Geol. Surv., Circ. 398.
- Mann, J. E., and H. E. Schlichting, Jr., 1967. Benthic algae of selected thermal springs in Yellowstone National Park. Trans. Am. Micros. Soc., 86(1): 2-9.
- Mann, K. H., 1958. Occurrence of an exotic oligochaete Branchiura sowerbyi Beddard, 1892, in the River Thames. Nature, London, 182: 732.
- Mann, K. H., 1965. Heated effluents and their effects on the invertebrate fauna of rivers. Proc. Soc. Wat. Treat. Exam., 14: 45-53.
- Mantelman, I. I., 1960. Distribution of the young of certain species of fish in temperature gradients. Translation Ser. No. 257, Fish. Resch. Bd. Can., Bio. Sta. Nanaimo, B. C., 67 pp. Spo. Fish. Abstr., 6(3), Abstr. No. 4381, 1961.

Mariner, L. T., and W. A. Hunsucker, 1959. Ocean cooling water systems for two thermal plants. Jour. of the Power Div., ASCE, 85(P04), Proc. Paper 2133.

Markowski, S., 1959. The cooling water of power stations: a new factor in the environment of marine and freshwater invertebrates. Jour. of Animal Ecol., 28(2): 243-258.

Markowski, S., 1960. Observations on the response of some benthonic organisms to power station cooling water. Jour. Animal Ecol., 29(2): 349-357. Spo. Fish. Abstr., 6(3), Abstr. No. 4263, 1961.

Markus, H. C., 1932. The extent to which temperature changes influence food consumption in largemouth bass (Huro floridana). Trans. Am. Fish. Soc., 62: 202-210.

Marrow, J. E., Jr., and A. Mauro, 1950. Body temperature of some marine fishes. Copeia, 2: 108-116.

Martin, N. V., 1951. A study of the lake trout Salvelinus namaycush, in two Algonquin Park, Ontario, lakes. Trans. Am. Fish. Soc., 81: 112-137.

Matsui, K., 1940. Temperature and heartbeat in a fish embryo, Oryzias latipes. I. The relationship of temperature coefficient of heartbeat to embryonic age. Sci. Rep. Tokyo Bun. Daig. 5B, 81-83: 39-51.

Matsui, K., 1943. Temperature and heart beat in a fish embryo, Oryzeas latipes, IV. The arrest of heartbeat by heat. Sci. Rep. Tokyo, Bun. Daig. 6B 95-98: 129-138.

Mattley, C. M., 1931. The effect of temperature on the number of scales in trout. Science, 74: 316.

Mayer, A. G., 1914. The effects of temperature upon tropical marine animals. Pap. Tortugas Lab. 6: 1-24.

McCauley, R. W., 1963. Lethal temperatures of the developmental stages of the sea lamprey, Petromyzon marinus L. Jour. Fish. Resch. Bd. Can., 20: 483. Jour. Water Poll. Control Federation, 36(7): 796, 1964.

McCay, C. M., L. A. Maynard, J. W. Titcomb, and M. F. Crowell, 1930. Influence of water temperature upon the growth and reproduction of brook trout. Ecol., 11: 30-34.

McCombie, A. M., 1960. Actions and interactions of temperature, light intensity and nutrient concentration on the growth of the green alga, Chlamydomonas reinhardi Dangeard. Jour. Fish. Resch. Bd. Can., 17(6): 871.

McEwen, G. F., 1929. A mathematical theory of the vertical distribution of temperature and salinity in water under the action of radiation, conduction, evaporation, and mixing due to the resulting convection. Bull., Scripps Institute of Oceano., LaJolla, Calif., Technical Ser. 2, No. 6, Univ. of Calif. Press.

McFarland, W. M., 1955. Upper lethal temperature in the salamander, Taricheer torosa, as a function of acclimation. Copeia, 3: 191-194.

McGraw, B. M., 1959. Ecology of the snail L. humilis Say. Trans. Am. Micro. Soc., 78(1): 101-121.

McKee, J. E., and H. W. Wolf, 1963. Water quality criteria. State Water Quality Contr. Bd., Calif., Publ. No. 3, 548 pp.

McLaren, I. A., 1963. Effects of temperature on growth of zooplankton, and the adaptive value of vertical migration. Jour. Fish. Resch. Bd. Can., 20: 684-727. Spo. Fish. Abstr., 8(3), Abstr. No. 6016, 1963.

McLeese, D. W., 1956. Effects of temperature, salinity and oxygen on the survival of the American lobster. Jour. Fish. Resch. Bd. Can., 13(2): 247-272.

McLeese, D. W., and D. G. Wilder, 1958. The activity and catchability of the lobster (Homarus americanus) in relation to temperature. Jour. Fish. Resch. Bd. Can. 15(6): 1345-1354.

Medcof, J. C., 1962. Possible effects of Passamaquoddy power project on clams, scallops and shipworms in Canadian waters. Jour. Fish. Resch. Bd. Can., 19(5): 877-889.

Medcof, J. C., and A. W. H. Needler, 1941. The influence of temperature and salinity on the condition of oysters. Jour. Fish. Resch. Bd. Can., 5(3): 243.

Medlen, A. B., 1951. Preliminary observations on the effects of temperature and light upon reproduction of Cambusia affinis. Copeia, 2: 148-152.

Meek, E. M., 1922. Effect of temperature on growth of young blennies (Zoarces vivparous). Dove Mar. Lab. Rep. 11: 102.

Meisner, H. M., and C. P. Hickman, Jr., 1962. Effect of temperature and photoperiod on the serum proteins of rainbow trout, Salmo gairdneri. Can. Jour. Zool., 40(2): 127-130.

Mekhanik, F. Y., 1955. Growth and metabolism in the young of the Ladogian salmon and rainbow trout at various temperature. Zool. Zhur. (USSR), 35: 290. Spo. Fish. Abs., 3(4), 1958

Merriman, D., 1935. The effect of temperature on the development of the eggs and larvae of the cutthroat trout (Salmo clarki clarki) Richardson. Jour. Exp. Bio., 12: 297-305.

Messinger, H., 1963. Dissipation of heat from a thermally loaded stream. Professional Paper No. 104, Geol. Surv., U. S. Dept. of the Interior.

Meuwis, A. L., and M. J. Heuts, 1957. Temperature dependence of breathing rate of carp. Bio. Bull. 112(1): 97-107.

Mihursky, J. A., 1962. Patuxent River estuary study with special references to the effects of heated steam electric station condenser water upon estuarine ecology. Unpublished Rep. No. 63-6, 29 pp., Nat. Resources Institute of Maryland.

Mihursky, J. A., and A. J. McErlean, 1964. Progress report to Department of Water Resources -- Patuxent River study with special reference to the effects of heated steam electric station condenser water upon estuarine ecology. Chesapeake Bio. Lab. Ref. No. 64-28.

Mikheev, V. P., 1962. On the distribution of Dreissena polymorpha Pall. on the constructions of the Volga hydro-electric station named Lenin. Byulletin Instituta Biologii Vodokhranilischeh. Akad. Nauk. USSR, 12: 32-33.

Miller, W. T., 1956. Possible relationship of water temperatures with availability and year class size in the Pacific sardine. M.A. Thesis in Bio. Sci., Stanford Univ.

Molnar, G., and I. Tolg, 1962. Relation between water temperature and gastric digestion of largemouth bass (Micropterus salmoides Lacepede). Jour. Fish. Resch. Bd. Can., 19(6): 1005-1012.

Monin, A. S., 1959. On the similarity of turbulence in the presence of a mean vertical temperature gradient. Jour. of Geophysical Resch., 64(12), Dec.

Mookerjee, H. K., D. N. Ganguly, and S. N. S. Gupta, 1946. The optimum range of temperature of water for the fingerlings of major carps of India. Sci. and Cult., 11: 383. Water Poll. Abstr. 19.

Moore, E. W., 1958. Thermal pollution of streams. Ind. Engng. Chem., 50: 4 87A. Spo. Fish. Abstr., 4(2), Abstr. No. 2428, 1959.

Moore, H. B., and E. G. Corwin, 1956. The effects of temperature, illumination and pressure on the vertical distribution of zooplankton. Bull. Mar. Sci. Gulf and Caribbean, 6(4): 273-287.

- Moore, J. A., 1949. Patterns of evolution in the genus Rana. In: Genetics, Paleontol., and Evol. Jepson, C. T., E. Mayr, and G. G. Simpson, Eds. pp. 315-338. Princeton Univ. Press.
- Moore, W. G., 1942. Field studies on the oxygen requirements of certain freshwater fishes. *Ecol.*, 23: 319-329.
- Moore, W. G., 1955. Observations on heat death in the fairy shrimp, Streptocephalus seali. *Proc. Louisiana Acad. Sci.*, 18: 5-12.
- Morris, R. W., 1961. Distribution and temperature sensitivity of some eastern Pacific cottid fishes. *Physiol. Zool.* 34: 217-227.
- Morris, R. W., 1962. Body size and temperature sensitivity in the cichlid fish, Aequidens portalegrensis (Hensel). *Am. Nat.*, 96: 35-50.
- Morton, K. E., 1962. Experimental heating of pond water to start rainbow trout on a dry diet. *Prog. Fish. Cult.*, 24(2): 94-96.
- Moss, D. D., and D. C. Scott, 1964. Respiratory metabolism of fat and lean channel catfish. *Prog. Fish Cult.*, 26(1): 16-20.
- Mossman, W. H., and A. L. Pacheco, 1957. Shad catches and water temperatures in Virginia. *Jour. Wildl. Mgmt.*, 21(3): 351-352.
- Murray, M. J., 1938. Survey of some northern Indiana streams with special reference to trout production. *Invest. Ind. Lakes*, 1: 79-99.
- Musacchia, J., and M. R. Clark, 1957. Effects of elevated temperatures on tissue chemistry of the Arctic sculpin, Myoxocephalus quarcornis. *Physiol. Zool.*, 30(1): 12-17.
- Nakai, N., 1927. On the influence of water temperature upon the development of the eggs of Leuciscus hakuensis Gunther. *Jour. Imp. Fish. Inst. Tokyo.*, 22: 73-85.
- Nakai, N., 1928. On the influence of the water temperature upon the hatching of eggs of Hypomessus alidus Pallas. *Jour. Imp. Fish. Inst. Tokyo*, 23: 124-127.
- Nakai, N., 1928. On the influence of temperature upon the hatching of eggs of Plecoglossus altinensis (TandS). *Jour. Imp. Fish. Inst. Tokyo*, 24: 28-37.
- Nakano, T., 1961. Studies on the physiological chemistry of phosphorus compounds in fish muscle V Quantitative difference of phosphorus compounds in muscle of fish reared in different water temperatures. *Bull. Jap. Soc. Sci. Fish.*, 24(4): 357-360. *Bio. Abstr.*, 41(3), 1962.

Naumann, E., 1927. Critique of the plankton concept. Arkiv. Bot. 21 A (3): 11-18.

Naylor, E., 1959. The fauna of a warm dock. Proc. XVth Int. Congr. Zool. Sect., 3, 259-262.

Naylor, E., 1963. Temperature relationships of the locomotor rhythm of Carcinus. Jour. Exp. Bio., 40: 669-679.

Naylor, E., 1965. Biological effects of a heated effluent in docks at Swansea, S. Wales. Proc. Zool. Soc., London, 144: 253-268. Water Poll. Abstr., 39(6), Abstr. No. 997, 1966.

Naylor, E., 1965. Effects of heated effluents upon marine and estuarine organisms. In: Advances in Marine Biology, Vol. 3, Academic Press, New York, pp. 63-103.

Needham, P. R., 1961. Observations on the natural spawning of Eastern brook trout. Cal. Fish and Game, 47(1): 27-40. Spo. Fish. Abs., 6(2), 1961.

Needham, P. R., and A. C. Jones, 1959. Flow, temperature, solar radiation, and ice in relation to activities of fishes in Sagehen Creek, California. Ecol., 40(3): 465-474.

Neess, J. C., and W. W. Bunge, 1957. An unpublished manuscript of E. A. Birge on the temperature of Lake Mendota. Part II. Trans. Wis. Acad. Sci., Arts and Letters, 46: 31-89.

Newcombe, C. L., and J. V. Slater, 1948. The occurrence of temperatures unusual to American lakes. Science, Vol. 108, Oct. 8.

Nicholls, J. W. V., 1931. The influence of temperature on digestion in Fundulus heteroclitus. Contr. Can. Bio., 7: 45-55.

Nielson, E. T., 1938. Thermoelectric measurement of the body temperature of mice and fishes. Acta. Med. Scand. 90(Suppl.): 169-189.

Noland, L. E., and E. Reichel, 1943. Life cycle of Lymnaea stagnalis completed at room temperature without access to air. Nautilus, 57(1): 8-13.

Nordlie, F. G., 1966. Thermal acclimation and peptic digestive capacity in the black bullhead Ictalurus melas (Raf.). Am. Midl. Nat., 75(2): 416-424.

Odum, E. P., 1959. Fundamentals of ecology 2nd ed. W. B. Saunders and Company, Philadelphia.

Ogilvie, D. M., and J. M. Anderson, 1965. Effect of DDT on temperature selection by young Atlantic salmon, Salmo salar. Jour. Fish. Resch. Bd. Can., 22(2): 503-512. Bio. Abstr., 46(17), Abstr. No. 74008, 1965.

Olson, P. A., 1958. Temperature tolerance of eggs and young of Columbia River fish. Hanford Bio. Res. Ann. Rep. for 1957. Off. of Tech. Ser., U. S. Dept. Commerce, Washington, D. C., p. 211.

Olson, P. A., and R. F. Foster, 1957. Temperature tolerance of eggs and young of Columbia River chinook salmon. Trans. Am. Fish. Soc., 85: 203.

Ordal, E. J., and R. E. Pacha, 1963. The effects of temperature on disease in fish. In: Water Temperature Influences, Effects and Control. Proc. of the 12th Pacific Northwest Symposium on Water Poll. Resch., Dept. of Health, Education and Welfare, Publ. Health Serv., Pacific Northwest Water Lab., pp. 39-56.

Orr, P. R., 1955. Heat death I Time-temperature relationships in marine animals. Physiol. Zool., 28(4): 290-294. Spo. Fish. Abstr., 2(2), Abstr. No. 788, 1957.

ORSANCO, 1956. Aquatic life water quality criteria, second progress report. Sewage and Indus. Wastes, 28(5): 678-690.

Orska, J., 1956. The influence of temperature on the development of the skeleton in teleosts. Zool. Polaniae, 7(3): 272-325.

Orton, J. H., 1920. Sea temperature, breeding and distribution of marine animals. Jour. Mar. Bio. Asso. Unit. King. 12: 339-366.

Palayer, P., 1963. Influence de la temperature sur la morphologie du pancreas de l'anguille (Anguilla anguilla L.). Bull. Mus. Hist. Nat., Paris, 35(2): 247-252.

Paloheimo, J. E., and L. M. Dickie, 1966. Food and growth of fishes. II. Effects of food and temperature on the relation between metabolism and body weight. Jour. Fish. Resch. Bd. Can., 23(6): 869-908.

Pannell, J. P. M., A. E. Johnson, and J. E. G. Raymont, 1962. An investigation into the effects of warmed water from Marchwood Power Station into Southampton water. Proc. Inst. Civil Engng. 23: 35-62.

Parker, B. W., 1965. Thermal loading. In: Minutes, Third Conf. on Patuxent Estuary Studies, Nov. 13-14, 1964. Chesapeake Bio. Lab. Ref. No. 6 -23.

Parry, G., 1961. Osmotic and ionic changes in blood and muscle of migrating salmonids. Jour. Exp. Bio., 38(2): 411-427. Spo. Fish. Abstr., 6(4), 1961.

- Parson, J. W., and E. Crittenden, 1959. Growth of redeye bass in Chipola River, Florida. Trans. Am. Fish. Soc., 88(3): 191-192.
- Patalas, K., 1960. Thermal and oxygen conditions and transparency of water in forty-four lakes of Wegoszewo District. Roczn. Nauk Rolniczych, Ser. B Zootech., 77(1): 105-222. Spo. Fish. Abstr., 7(4), 1962.
- Patel, B., 1959. The influence of temperature on the reproduction and moulting of Lepas anatifera L. under laboratory conditions. Jour. Mar. Bio. Asso. Unit. King., 38: 589-597.
- Patel, B., and D. J. Crisp, 1960. The influence of temperature on the breeding and moulting activities of some warm water species of operculate barnacles. Jour. Mar. Bio. Asso. Unit. King., 39: 667-680.
- Patrick, R., 1949. Proposed biological measure of stream conditions, based on a survey of the Conestoga Basin, Lancaster County, Pa. Proc. Acad. Nat. Sci. Phila., CI: 277-341.
- Patrick, R., M. H. Hahn, and J. H. Wallace, 1954. A new method for determining the pattern of diatom flora. Norulae Natulae, Acad. Nat. Sci. of Phila. 259: 1-12.
- Pearce, D. W., 1965. Plankton of an artificially heated dock. Ann. Rep. Challenger Soc., 3(16): 24-25.
- Pearse, A. S., and F. G. Hall. 1928. Homoithermism. The origin of warm blooded vertebrates. John Wiley and Sons, New York, 119 pp.
- Pearson, B. E., 1952. The behaviour of a sample of hybrid trout (Salvelinus fontinalis x Cristivomer namaycush) in a vertical temperature gradient. Ontario Fish. Resch. Lab. Library, Toronto, 24 pp. M.S.
- Pegel', V. A., 1959. Mechanism of adaptation by fishes to the temperature factor. In: Bio. fundamentals of fishing industry. Tomsk: Tomskii Univ. 1959: 135-142. Referat. Zhur., Bio. 1961, No. 4D473. Bio. Abstr., 40(5), No. 18899, 1962.
- Peiss, C. M., and J. Field, 1950. The respiratory metabolism of excised tissues of warm and cold adapted fishes. Bio. Bull., 99(2): 213-224.
- Penfound, W. T., 1956. Primary production of vascular aquatic plants. Limnol. and Oceano., 1(2): 92-101.
- Petryaev, E. D., and N. T. Tayurskaya, 1945. Effectiveness of water chlorination at low temperatures. Gigiena, USSR, 10(9): 52. Water Poll. Abstr., Vol. 20, May 1947.

- Pfitzer, D. W., 1954. Investigations of water below storage reservoirs in Tennessee. Trans. 19th N. A. Wildl. Conf., pp. 271-282.
- Phillips, A. M., Jr., 1961. Effect of water temperature and diet on the blood glucose of brook trout. Prog. Fish. Cult., 23(2): 66-69.
- Phillips, A. M., Jr., 1962. Effect of diet and water temperature on the blood phosphorus of brook trout. Prog. Fish. Cult., 24(1): 22-25.
- Piavis, G. W., 1961. Embryological stages in the sea lamprey and effects of temperature on development. Bureau of Commercial Fisheries, U. S. Fish and Wildl. Serv., Fish. Bull. 182, 61: 111-143.
- Pickard, G. L., and R. W. Trites, 1957. Freshwater transport determination from the heat budget with application to British Columbia inlets. Jour. Fish. Resch. Bd. Can., 14(4): 605-616.
- Picton, W. L., 1956. Water use in the United States, 1900-1975. Business Serv. Bull. 136, Water and Sewage Indus. and Util. Div., U. S. Dept. of Commerce.
- Pitkow, R. B., 1960. Cold death in the guppy. Bio. Bull., 199(2): 231-245.
- Pitt, T. K., E. T. Garside, and R. L. Hepburn, 1956. Temperature selection of the carp (Cyprinus carpio L.). Can. Jour. Zool., 34: 555-557.
- Podoliak, H. A., 1961. Relation between water temperature and metabolism of dietary phosphorus by fingerling brook trout. Trans. Am. Fish. Soc., 90: 398-403.
- Powers, E. B., et al, 1932. The relation of respiration of fishes to environment. Ecol. Monog., 2(4): 385-473.
- Powers, E. B., 1937. Factors involved in the sudden mortality of fishes. Trans. Am. Fish. Soc., 67: 270-280.
- Pratt, D. M., 1943. Analysis of population development in Daphnia at different temperatures. Bio. Bull., 85(2): 116-140.
- Pratt, D. M., and D. A. Campbell, 1956. Environmental factors affecting growth in Venus mercenaria. Limnol. and Oceano. 1(1): 2.
- Precht, H., J. Christophersen, and H. Hensel, 1955. Temperatur und leben. Springer Verlag, Berlin, Germany, 514 pp.
- Price, J. W., 1940. Time-temperature relations in the incubation of the white fish, Coregonius clupeoformis (Mitchill). Jour. Gen. Physiol. 23: 449-468.

Pritchard, D. W., 1965. Prediction of excess temperature distribution. In: Minutes, Third Ann. Conf. on the Patuxent Estuary Studies, Nov. 13-14, 1964. Chesapeake Bio. Lab. Ref. No. 65-23.

Pritchard, D. W., and H. H. Carter, 1965. On the prediction of the distribution of excess temperature from a heated discharge in an estuary. Tech. Rep. 33, Ref. 65-1, Chesapeake Bay Institute, The Johns Hopkins Univ., 45 pp.

Privol'nev, T. I., and N. V. Koroleva, 1953. Critical content of oxygen in water for fish at various temperatures according to seasons. *Compt. Rend. Acad. Sci. USSR*, 89: 175. *Sewage and Ind. Wastes*, 27(6): 666, 1955.

Privol'nev, T. I., 1963. Threshold concentrations of oxygen in water for fish at various temperatures. *Dokl. Akad. SSR*, 151(2): 439. *Jour. Water Poll. Control Federation*, 36(7): 795, 1964.

Prosser, C. L. Physiological variations in animals. *Bio. Rev.*, 30(3): 229-262.

Prosser, C. L., 1957. The species problem from the viewpoint of a physiologist. In: *The species problem*. Am. Asso. Adv. Sci., Washington, D. C., pp. 339-369.

Prosser, C. L., 1963. Metabolic and nervous acclimation of fish to cold and heat. *Abstr. Int. Symp. Cytoecol. Moscow*: 49-50.

Prosser, C. L., F. A. Brown, D. W. Bishop, T. L. John, and V. J. Wulff, 1950. *Comparative animal physiology*. W. B. Saunders Co., Phila., Pa., 888 pp.

Provasoli, L., 1958. Nutrition and ecology of protozoa and algae. *Ann. Rev. of Microbio.*, 12: 279-303.

Purkett, C. A., Jr., 1950. Growth rate of trout in relation to elevation and temperature. *Trans. Am. Fish Soc.*, 80: 251-259.

Raffy, A., 1952. Influence des variations de la temperature sur l'osmoregulation des petites carpes en eau douce et en eau salée. *Compte Rendus Soc. de Bio.* 146: 908-910.

Raney, E. C., 1966. Bibliography. Heated discharges and effects on aquatic life with emphasis on fishes. Division of Biological Sciences, Section of Ecology and Systematics, Fernow Hall, Cornell Univ., Ithaca, New York. Oct. 19, 1966, 34 pp. (mimeo.).

Rao, K. P., 1953. Shell weight as a function of intertidal height in a littoral population of pelecypods. *Experimentia*, 9: 465.

Rao, K. P., 1953. Threshold concentrations of oxygen in water for fish at various temperatures. Dokl. Akad. SSR, 151(2): 439. Jour. Water Poll. Control Federation, 36(7): 795, 1964.

Rao, K. P., 1962. Physiology of acclimation to low temperature in poikilotherms. Science, 137(3531): 682-683.

Rao, K. P., and T. H. Bullock, 1954. Q as a function of size and habitat temperature in poikilotherms. Am. Nat., 88: 33-44.

Raphael, J. M., 1962. Prediction of temperature in rivers and reservoirs. Jour. of the Power Div., ASCE, 88(P02), Proc. Paper 3200.

Rawson, D. S., 1956. Life history and ecology of the yellow walleye, Stizostedion vitreum, in Lac La Ronge, Saskatchewan. Trans. Am. Fish. Soc., 86: 15-37.

Rawson, D. S., 1960. A limnological comparison of twelve large lakes in northern Saskatchewan. Limno. and Oceano., 5(2): 195-211.

Read, K. R. H., 1962. Respiration of the bivalved molluscs Mytilus edulis L. and Brachidontes demissus plicatulus Lamarck as a function of size and temperature. Comp. Biochem. and Physiol., 7(1-2): 89-101.

Rees, W. J., 1954. Biology of pollution. Jour. Inst. Bio., 1: 29-30.

Reid, G. W., and R. Zaltman, 1964. Thermal pollution. Prepr. 4th Ind. Wat. Waste Conf., Tex. Wat. Poll. Control Asso., A.2-A.14.

Reimers, N., 1963. Body conditions, water temperature, and over-winter survival of hatchery-reared trout in Convict Creek, California. Trans. Am. Fish. Soc., 92(1): 39-46.

Reish, D. J., 1961. The relationship of temperature and dissolved oxygen to the seasonal settlement of the polychaetous annelid Hydroides norvegica (Gunnerus). Bull. S. Calif. Acad. Sci. 60: 1-11.

Renn, C. E., 1957. Warm-water effects on municipal supplies. Jour. Am. Water Works Asso, 49: 405.

Ricker, W. E., 1934. An ecological classification of certain Ontario streams. Univ. Tor. Stud. Bio. Ser. 37, Publ. Ont. Fish. Resch. Lab., 49: 1-114.

Ricker, W. E., 1946. Industrial and domestic wastes in relation to aquatic life of Indiana streams. Purdue Univ. Engng. Bull. No. 60. Proc. of the 2nd Indus. Wastes Conf., Jan. 10.

Roberts, J. L., 1953. Studies on thermal acclimatisation in the lined shore crab, Pachygrapsus crassipes Randall. Proc. XIXth int. Physiol. Congr., p. 706.

- Roberts, J. L., 1957. Thermal acclimation of metabolism in the crab, Pachygrapsus crassipes Randall. II Mechanisms and the influence of season and latitude. *Physiol. Zool.* 30: 242-255.
- Rogers, C. G., and E. M. Lewis, 1916. The relation of the body temperature of certain cold-blooded animals to that of their environment. *Bio. Bull.*, 31: 1-15.
- Roots, B. I., and C. L. Prosser, 1962. Temperature acclimation and the nervous system in fish. *Jour. Exp. Bio.*, 39: 617-629.
- Ross, F. F., 1958. The operation of thermal power stations in relation to streams. The Inst. of Sew. Purif., Ann. Conf., Southport, June 24-25th. Conf. Pap. No. 7, 10 Cromwell Pl., South Kensington, London, S.W. 7.
- Ross, F. F., 1959. The operation of thermal power station in relation to streams. *Jour. Inst. Sew. Purif.*, No. 1: 16-26.
- Rozin, P. M., and J. Mayer, 1961. Thermal reinforcement and thermoregulatory behavior in the goldfish, Carassius auratus. *Science*, 134(3483): 942-943.
- Rubin, M. A., 1935. Thermal reception in fishes. *Jour. Gen. Physiol.*, 18: 643-647.
- Ruhr, C. E., 1954. Smallmouth stream management studied. *Tenn. Conserv.* 20(2): 10-13. *Spo. Fish. Abstr.*, 1(2), Abstr. No. 275, 1955.
- Runnstrom, S., 1928. Über die thermopathie der fortpflanzung und entwicklung mariner tiere in beziehung zu ihrer geographischen verbreitung. *Bergens Mus. Arb. Naturv rekke*, 1927, No. 2, 66 pp.
- Rutkovskiy, V. I., 1962. Distribution of temperature along the Upper Volga reservoirs in the period of intensive heating. *Byulletin Instituta Biologii Vodokhranilischeh, Akad. Nauk. USSR* 13: 48-51.
- Sand, A., 1938. The function of the ampollae of Lorenzii with some observations on the effect of temperature on sensory rhythms. *Proc. Roy. Soc. London* 125B, 841: 524-553.
- Saur, J. F. T., and E. R. Anderson, 1956. The heat budget of a body of water of varying volume. *Limnol. and Oceano.*, 1(4): 247-251.
- Sayle, M. H., 1928. Factors influencing the rate of metabolism of Aeschna umbrosa nymphs. *Bio. Bull.*, 54: 212.
- Schaperclaus, W., 1927. The acidity of fresh water and its relation to fish. *Sitzungsker ges Naturfarsch Freunde. Berlin* (1/3): 1-9.

- Scholander, P. F., W. Flagg, V. Walter, and L. Irving, 1953. Climatic adaptations in arctic and tropical poikilotherms. *Physical Zool.* 26: 67-92.
- Schrader, T., 1958. Thermal conditions in impounding reservoirs. *Limnol. Sch. Reihe.*, 5(21): 68-88. *Spo. Fish. Abstr.*, 5(3), 1960.
- Schultz, L. P., 1927. Temperature controlled variation in the golden shiner, Notemigonis crysoleus. *Pap. Mich. Acad.*, Ann Arbor, 7: 417-432.
- Schwartz, F. J., 1964. Effects of winter water condition on fifteen species of captive marine fishes. *Am. Midl. Nat.*, 71: 434-444. *Bio. Abstr.*, 45: 85134.
- Scott, D.C., 1958. Progress report on dissolved oxygen tolerance of bluegills at 25°C and 35°C. *Tech. Bull. No. 111*, Nat. Council for Stream Improvement.
- Scott, D. P., 1964. Thermal resistance of pike (Esox lucius L.), muskellunge (E. masquinongy Mitchill), and their F₁ hybrid. *Jour. Fish. Resch. Bd. Can.*, 21(5): 1043-1049.
- Segal, E., 1956. Microgeographic variation as thermal acclimation in an intertidal mollusc. *Bio. Bull. Woods Hole*, 111: 129-152.
- Segal, E., K. P. Rao, and T. W. James, 1953. Rate of activity as a function of intertidal height within populations of some littoral molluscs. *Nature (London)*, 172: 1108.
- Sette, A. E., 1950. Biology of the Atlantic mackerel (Scomber scombris) of North America, Part II. Migrations and habitats. *Fish. Bull. U. S.*, 49: 251-258.
- Seymour, A. H., 1956. Effects of temperature upon young chinook salmon. Ph.D. Thesis, Univ. Wash., Seattle, 127 pp. *Diss. Abstr.* 16(11): 2249. *Spo. Fish. Abstr.*, 2(3), Abstr. No. 953, 1957.
- Seymour, A., 1959. Effects of temperature upon the formation of vertebrate and fin rays in young chinook salmon. *Trans. Am. Fish. Soc.*, 88: 58-69.
- Sharpe, F. P., 1961. A mid-summer plant of rainbow trout in a stratified Tennessee impoundment. *Prog. Fish. Cult.*, 23(4): 152-155.
- Sheridan, W. L., 1961. Temperature relationships in a pink salmon stream in Alaska. *Ecol.*, 42(1): 91-98.
- Shaw, P. A., 1946. Oxygen consumption of trout and salmon. *Calif. Fish and Game*, 32(1): 3-12.

- Shetter, D. S., and M. J. Whalls, 1955. Effect of impoundment on water temperatures of Fuller Creek, Montmorency County, Michigan. Jour. Wildl. Mngmt., 19(1): 47-54. Spo. Fish. Abstr., 1(4), 1956.
- Shultz, L. P., 1927. Temperature controlled variation in the golden shiner, Notemigonus crysoleucus. Pap. Mich. Acad. 7: 417-432.
- Simpson, A. C., 1953. Some observations on the mortality of fish and the distribution of plankton in the southern North Sea during the cold winter, 1946-1947. Jour. Cons. Mt. Explor. Mer., 19: 150-177.
- Simpson, R. W., 1960. Urban redevelopment and thermal discharge. Wat. and Sewage Works, Nov., 1960: 448-449.
- Slack, K. V., and F. E. Clarke, 1965. Patterns of dissolved oxygen in a thermally loaded reach of the Susquehanna River, Pennsylvania. Prof. Paper U. S. Geol. Surv., 525-C: 193-195. Water Poll. Abstr., 39(7), Abstr. No. 1210, 1966.
- Smith, D. C., 1928. The effect of temperature on the melanophores of fishes. Jour. Exp. Zool., 52: 183-234.
- Smith, D. C., 1931. The effect of temperature changes upon the pulsations of isolated scale melanophores of Fundulus heteroclitus. Bio. Bull. 60: 269-287.
- Smith, L. L., et al., 1955. Stream pollution. Aquatic life water quality criteria. First Prog. Rep. Aquatic Life Advisory Comm. of the Ohio River Valley Water Sanitation Comm., Sewage and Indus. Wastes, 27(3): 321-331.
- Smith, L. L., et al., 1956. Stream pollution. Aquatic life water quality criteria. Second Prog. Rep. Aquatic Life Advisory Comm. of the Ohio River Valley Water Sanitation Comm. Sewage and Indus. Wastes, 28(5): 678-690.
- Somers, G. N., and A. L. DeVries, 1967. Temperature tolerance of some antarctic fishes. Science, 156: 257-258, April 14.
- Southgate, B. A., 1951. Pollution of streams: some notes on recent research. Inst. Civ. Engng., Publ. Health Paper No. 1.
- Southward, A. J., 1958. Note on the temperature tolerances of some intertidal animals in relation to environmental temperatures and geographical distribution. Jour. Mar. Bio. Asso. Unit. King., 37: 49-66.
- Spaas, J. T., 1960. Contribution to the comparative physiology and genetics of the European Salmonidae III Temperature resistance at different ages. Hydrobiologia, 15: 78-88.

Spector, W. S., 1956. Handbook of biological data. W. B. Saunders Co., 584 pp.

Spencer, R. W., and J. Bruce, 1960. Cooling water for steam electric stations on tidewater. Jour. of the Power Div., ASCE, 86(P03), Proc. Paper 2503, June.

Spoor, W. A., 1951. Temperature and the erythrocyte count of goldfish. Fed. Proc., 10: 131.

Sprague, J. B., 1963. Resistance of four freshwater crustaceans to lethal high temperature and low oxygen. Jour. Fish. Resch. Bd. Can., 20: 387.

Stander, G. J., 1952. Quality requirements of water for the maintenance of aquatic flora and fauna and for recreational purposes. So. Afr. Ind. Chem., 6: 118.

Stangenberg, M., 1965. Natural effects of draining warm waters off into the rivers. Gaz Woda Tech. Sanit., 39(3): 86-89. Chem. Abstr. 63, Abstr. No. 4007, 1965. Water Poll Abstr., 39(5), Abstr. No. 793, 1965.

Stangenberg, M., and M. Z. Pawlaczyk, 1961. The influence of a warm-water influx from a power station upon the formation of biocenotic communities in a river. Nauk. Pol. Wr., Wroclaw, No. 40, Inzyn. Sanit. I: 67-106. Water Poll. Abstr., 35(3), Abstr. No. 579, 1962.

Stankiewicz, E. J., 1958. Water supply to thermal power plants. Jour. of the Power Div., ASCE, 84(P06), Proc. Paper 1889, Dec.

Stevenson, J. H., and A. A. Hulsey, 1961. Vertical distribution of dissolved oxygen and water temperature in Lake Hamilton with special reference to suitable rainbow trout habitat. Proc. 15th Ann. Conf., Southeastern Asso. Game and Fish Comm., pp. 245-255. Spo. Fish. Abstr., 8(1), 1963.

Stoltenberg, D. H., 1965. Effects of temperature on the deoxygenation of a polluted estuary. Jour. Water Poll. Control Federation, Vol. 37, Dec.

Storey, M., 1937. The relation between normal range and mortality of fishes due to cold at Sanibel Island, Florida. Ecol., 18: 10-26.

- Strandberg, C. H., 1962. Dispersion and diffusion of heated coolant water. Gen. Resch. and Development, Inc., 112 Wayne St., Arlington, Va.
- Strandberg, C. H., 1962. Analysis of thermal pollution from the air. Proc. of 17th Purdue Wastes Conf., Purdue Univ., W. Lafayette, Ind.
- Strawn, K., 1961. Growth of largemouth bass fry at various temperatures. Trans. Am. Fish. Soc., 90(3): 334-335. Water Poll. Abstr., 35(3), Abstr. No. 580, 1962.
- Sullivan, C. M., 1949. Aspects of the physiology of temperature selection in speckled trout (Salvelinus fontinalis). Univ. Toronto, Dept. Zool., Ph.D. Thesis.
- Sullivan, C. M., 1954. Temperature reception and responses in fish. Jour. Fish. Resch. Bd. Can., 11(2): 153-170.
- Sullivan, C. M., and K. C. Fisher, 1953. Seasonal fluctuations in the selected temperature of speckled trout, Salvelinus fontinalis (Mitchill). Jour. Fish. Resch. Bd. Can., 10: 187-195.
- Sumner, F. B., and P. Doudoroff, 1938. Some experiments upon temperature acclimatization and respiratory metabolism in fishes. Bio. Bull., 74: 403-429.
- Sumner, F. B., and A. N. Lanahanm, 1942. Studies of the respiratory metabolism of warm and cool spring fishes. Bio. Bull., 82: 313-327.
- Sumner, F. B., and M. C. Sargent, 1940. Some observations on the physiology of warm spring fishes. Ecol., 21: 45-54.
- Sumner, F. B., and N. A. Wells, 1935. Some relations between respiratory metabolism in fishes and susceptibility to certain anaesthetics and lethal agents. Bio. Bull., 69(3): 368-378.
- Swain, A., and C. F. Newman, 1957. Hydrographical survey of the River Usk. Fish. Invest. Ser. I, 6(1).
- Swarup, H., 1958. Abnormal development in the temperature-treated eggs of Gasterosteus aculeatus (L.) I Cleavage abnormalities. Jour. Zool. Soc. India, 10: 108-113.
- Swarup, H., 1959. Abnormal development in the temperature-treated eggs of Gasterosteus aculeatus (L.) II Gastrulation abnormalities. Jour. Zool. Soc. India, 11: 1-6.
- Swarup, H., 1959. Abnormal developemtn in the temperature-treated eggs of Gasterosteus aculeatus (L.) III Twinning. Jour. Zool. Soc. India, 11: 7-10.

Swift, D. R., 1959. Seasonal variation in the activity of the thyroid gland of yearling brown trout, Salmo trutta L. Jour. Exp. Bio., 36(1): 120-125. Spo. Fish. Abstr., 4(2), 1959.

Sylvester, R. O., 1960. Some influences of multipurpose water usage on water quality. 7th Tech. Mtg., Inter. Union Cons. Nature and Nat. Resources, Theme I (4): 215-235. Spo. Fish. Abstr., 5(4), 1960.

Sylvester, R. O., 1963. Effects of water uses and impoundments on water temperature. Water temperature influences, effects, and control. U. S. Dept. of Health, Education and Welfare, Publ. Health Serv., Pacific Northwest Water Lab., Corvallis, Ore., pp. 6-27.

Tabata, S., 1958. Heat budget of the water in the vicinity of Triple Island, British Columbia. Jour. Fish. Resch. Bd. Can., 15(3), 1958.

Tahatz, M. E., 1961. Tolerance of striped bass and American shad to changes of temperature and salinity. U. S. Fish and Wildl. Serv., Spec. Sci. Rep., Fish. No. 388, 8 pp.

Tarzwel, C. M., 1957. Water quality criteria for aquatic life. In: Biological Problems in Water Pollution. Robert A. Taft Sanit. Engng. Center, Cincinnati, Ohio, pp. 246-272.

Tarzwel, C. M., 1958. Dissolved oxygen requirements for fishes. In: Oxygen Relationships in Streams. Proc. of Seminar, Cincinnati, Ohio, Oct. 30 - Nov. 1, 1957. Publ. Health Serv. Publ. No. W58-2.

Tarzwel, C. M., 1960. Effects and control of stream pollution. Soil and Water Conserv., Publ. Tech. Mtg. of LVCN, FAO, Sept. 1958.

Tarzwel, C. M. (Ed.), 1960. Biological problems in water pollution. Second Seminar 1959, Robert A. Taft Sanit. Engng. Center, Cincinnati, Ohio. Publ. Health Serv. Publ. No. W60-3.

Tarzwel, C. M. (Ed.), 1965. Biological problems in water pollution. Third Seminar 1962, Robert A. Taft Sanit. Engng. Center, Cincinnati, Ohio. Publ. Health Serv. Publ. No. 999-WP-25, 434 pp.

Tarzwel, C. M., and A. R. Gaufin, 1958. Some important biological effects of pollution often disregarded in stream surveys. Proc. 8th Purdue Ind. Waste Conf.

Tauti, M., 1927. On the influences of temperature and salinity upon the rate of development of fish eggs. Jour. Imp. Fish. Inst. Tokyo, 23: 31-37.

Tauti, M., 1928. On the influence of temperature of water upon the hatch rate and the hatching days of fish eggs. Jour. Imp. Fish. Inst. Tokyo, 24(1): 13-18.

- Teter, H. E., 1960. The bottom fauna of Lake Huron. Trans. Am. Fish. Soc., 89(2): 193-197.
- Thornton, F. E., 1932. The viscosity of the plasmogel of Amoeba proteus at different temperatures. Physiol. Zool., 5(2): 246-253.
- Threinen, C. W., 1958. Cause of mortality of a midsummer plant of rainbow trout in a southern Wisconsin lake, with notes on acclimation and lethal temperatures. Prog. Fish. Cult., 20: 27. Spo. Fish. Abstr., 3(2), Abstr. No. 1543, 1958.
- Tkachenko, N. I., 1960. Effect of temperature on the microflora of biofilters. Sb. Tr. Gos. Nauch.-Issled. Inst. Gidroliz. Sulf.-Spirt. Prom., 1960, 8: 252-262. Abstr. Bull. Inst. Pap. Chem., 1963, 33:676.
- Trembley, F. J., 1960. Research project on effects of condenser discharge water on aquatic life, progress report, 1956 to 1959. Inst. of Resch., Lehigh Univ., 154 pp. Water Poll. Abstr., 34(11), Abstr. No. 2157, 1961.
- Trembley, F. J., 1961. II Progress report, 1960. Lehigh Univ. Inst. of Resch., Bethlehem, Pa., Unpubl.
- Trembley, F. J., 1965. Effects of cooling water from steam. In: Biological Problems in Water Pollution (Tarzwell, C. M., Ed.). Publ. Health Serv. Publ. No. 999-WP-25, pp. 334-345.
- Trembley, L., 1962. Temperature de l'eau d'un lac et la migration de frai du Catostome, Catostomus c. commersoni. Nat. Canad., 89: 119-128.
- Truesdale, G. O., and K. G. VanDyke, 1958. Effect of temperature on aeration of flowing water. Water and Waste Treatment Jour., 7: 9-11.
- Tsukuda, H., 1960. Temperature adaptation in fishes IV Change in the heat and cold tolerances of the guppy in the process of temperature acclimatization. Jour. Inst. Polyt. Osaka Univ., 11D: 43-54.
- Tsukuda, H., 1960. Heat and cold tolerances in relation to body size in the guppy, Lebistes reticulatus. Jour. Inst. Polyt. Osaka Univ., 11D: 55-62.
- Tsukuda, H., 1961. Temperature acclimatization on different organization levels in fishes. Jour. Bio. Osaka City Univ. 12: 15-45. Bio. Abstr., 45(8), Abstr. No. 33313, 1964. Spo. Fish. Abstr. 10(1), Abstr. No. 7326, 1965.
- Tsukuda, H., and T. Katayama, 1957. Temperature adaptation in fishes I. The influences of rearing temperature on the temperature tolerance, growth rate and body form. Physiol. Ecol., Kyoto, 7: 113-122 (English summary). Spo. Fish. Abstr., 4(4), Abstr. No. 2815, 1959.

Tsukuda, H., and W. Ohsawa, 1959. The heat and cold coma temperatures in Lebistes reticulatus as determined by raising and lowering water temperature at constant rates. Jour. Inst., Polyt. Osaka Univ., 9D: 69-76.

Tucker, D. S., 1958. The distribution of some fresh water invertebrates in ponds in relation to annual fluctuations in the chemical composition of the water. Jour. Animal Ecol., 27: 105-123.

VanOosten, J., 1944. Factors affecting the growth rate of fish. Trans. North Am. Wildl. Conf., 9: 177-183.

Varley, M., 1961. Biological effects of power station discharge. Power Supplies and Water Resources Symposium, 1961. Reprint by Engineering.

Vaux, D., 1953. Hydrographical conditions in the southern North Sea during the cold winter of 1946-1947. Jour. Cons. Int. Explor. Mer. 19: 127-149.

Vellas, F., 1963. Influence de la temperature sur les activitises uricolytiques de quelques Teleosteens dulcicoles. C. R. Acad. Sci. Paris, 256: 4740-4743.

Velz, C. J., 1939. Deoxygenation and reoxygenation. Trans. Am. Soc. Civil Engng., 104: 560-578.

Velz, C. J., 1944. Factors influencing self-purification and their relation to pollution abatement. Sewage and Indus. Wastes, 19: 629-644.

Velz, C. J., and J. J. Gannon, 1960. Forecasting heat losses in ponds and streams. Sewage and Indus. Wastes, 32(4): 392. Spo. Fish. Abstr. 6(1), Abstr. No. 3917, 1961.

Verduin, J., 1960. Letter in science 131 p. 232, January 22, 1960.

Vernick, S. H., 1962. The effects of temperature, light and high frequency radio waves upon the embryonic development of Tilapia macrocephala (Bleeker). Dissertation Abstr., 23(4): 1174-1175. Bio. Abstr., 42(5), Abstr. No. 18251, 1962.

Vernon, H. M., 1899. The death-temperature of certain marine organisms. Jour. Physiol. 26: 131-136.

Verril, A. E., 1901. A remarkable instance of the death of fishes at Bermuda in 1901. Am. Jour. Sci. Ser. 4, 12: 88-95.

Viehl, K., 1950. The influence of temperature on biological decomposition in water and sludge with special reference to the effect of discharges of hot water on streams. Gesundheitsing, 71: 349. Water Poll. Abstr., 27(7), Abstr. 168, 1952.

- Vliet, R. V., 1957. Effect of heated condenser discharge water upon aquatic life. Am. Soc. Mech. Engrs. Paper No. 57-PWR-4, 10 pp. Water Poll. Abstr., 31(11), Abstr. No. 2242, 1958.
- Volpe, E. P., 1957. Embryonic temperature adaptations in highland Rana pipiens. Am. Nat. 91(860): 303-310.
- Waede, M., 1954. Osmotic, chemical and thermal resistance of sole (Pleuronectes platessa) and flounder (P. flesus). Kiel. Meereforsch, 10: 58. Sewage and Indus. Wastes, 27(6), 1955.
- Walford, L. A., 1938. Effect of currents on the distribution and survival of the eggs and larvae of the haddock (Melanogrammus aeglefinus) on Georges Bank. Bull. U. S. Bur. Fish., 49: 1-73.
- Wallace, N. M., 1955. The effect of temperature on the growth of some fresh water diatoms. Not. Nat. Acad. Nat. Sci., No. 280, 11 pp.
- Walshe, B. M., 1948. Oxygen requirements and thermal resistance of chironomid larvae from flowing and still water. Jour. Exp. Bio., 25: 35.
- Ware, G. C., 1958. Effect of temperature on the biological destruction of cyanide. Water and Waste Treatment Jour., 6: 537.
- Warinner, J. E., and M. L. Brehmer, 1964. The effects of thermal effluents on marine organisms. Proc. 19th Waste Conf. Purdue Univ., Engng. Extn. Ser. 117: 479-492. Water Poll. Abstr., 39(4), Abstr. No. 625, 1966.
- Warinner, J. E., and M. L. Brehmer, 1966. The effects of thermal effluents on marine organisms. International Jour. Air and Water Poll., 10(4): 277-289.
- Watts, R. L., and G. W. Harvey, 1946. Temperature of Kettle Creek and tributaries in relation to game fish. Bull. Penn. Agr. Exp. Sta., 481: 1-30.
- Weatherley, A. H., 1959. Some features of the biology of the tench Tinca tinca (L.) in Tasmania. Jour. Animal Ecol., 28: 73-87.
- Weatherly, A. H., 1963. Thermal stress and interrenal tissue in the perch Perca fluviatilis (L.). Proc. Zool. Soc. London, 141(3): 537-555. Spo. Fish. Abstr., 9(1), Abstr. No. 6314, 1964.
- Weatherly, A. H., 1963. Zoogeography of Perca fluviatilis (L.) and Perca flavescens (Mitchill) with special reference to the effects of high temperature. Proc. Zool. Soc. London, 141(3): 557-576. Spo. Fish. Abstr., 9(1), Abstr. No. 6315, 1964.

Webster, D. W., 1945. Relation of temperature to survival and incubation of the eggs of smallmouth bass (Micropterus dolomieu). Trans. Am. Fish. Soc. , 75: 43-47.

Wells, H. W., et al, 1961. Winter fish mortality in Pamlico Sound, North Carolina. Ecol., 42: 217-219.

Wells, M. M., 1914. Resistance and reactions of fishes to temperature. Trans. Illinois Acad. Sci., 7: 48-59.

Wells, N. A., 1935. Variations in the respiratory metabolism of the Pacific killifish, Fundulus parvipinnis, due to size, season, and continued constant temperature. Physiol. Zool., 8: 318-336.

Wells, N. A., 1935. Change in rate of respiratory metabolism in a teleost fish induced by acclimatization to high and low temperature. Bio. Bull. Woods Hole, 69: 361-367.

Whitemore, C. M., C. E. Warren, and P. Doudoroff, 1960. Avoidance reactions of salmonid and centrarchid fishes to low oxygen concentrations. Trans. Am. Fish. Soc., 89(1): 17-26.

Whitney, R. J., 1939. The thermal resistance of mayfly nymphs from ponds and streams. Jour. Exp. Bio. 16: 374-385.

Whitworth, W. R., and W. H. Irwin, 1964. Oxygen requirements of fishes in relation to exercise. Trans. Am. Fish. Soc., 93(2): 209-212. Spo. Fish. Abstr., 9(3), Abstr. No. 6772, 1964. Water Poll. Abstr., 39(6), Abstr. No. 991, 1966.

Wickwire, G. C., L. D. Sager, and W. E. Burge, 1929. Comparative effect of temperature on rate of pure chemical reactions and rate of sugar utilization by a plant and on cold blooded animal (Spirogyra and goldfish). Bot. Gaz., 88(4): 430-436.

Williams, A. E., and R. K. Burris, 1952. Nitrogen fixation by blue-green algae and their nitrogenous composition. Am. Jour. Bot., 39: 340-342.

Wingfield, C. A., 1939. The activity and metabolism of poikilothermal animals in different latitudes. IV. Proc. Zool. Soc. London, A109: 103-108.

Wingfield, C. A., 1940. The effect of certain environmental factors on the growth of brown trout (Salmo trutta). Jour. Exp. Bio., 17: 435-438.

Wohlschlag, D. E., 1960. Metabolism of an antarctic fish and the phenomenon of cold adaptation. Ecol., 41(2): 287-292. Spo. Fish. Abstr., 5(3), Abstr. No. 3418, 1960.

- Wood, A. H., 1932. The effect of temperature on the growth and respiration of fish embryos (Salmo fario) Jour. Exp. Bio., 9: 271-276.
- Woodward, D. R., 1957. Availability of water in the United States with special reference to industrial needs by 1980. Industrial College of the Armed Forces, Thesis No. 143.
- Woynarovich, E., 1961. Oxygen consumption of Dreissena polymorpha at different temperatures. Magy. Tud. Akad. Tihanyi. Bio. Kutatoint., Evkonye, 28: 211. Sewage and Indus. Wastes, 35(7); 1961.
- Wuhrmann, K., and H. Woker, 1955. Influence of temperature and oxygen tension on the toxicity of poisons to fish. Proc. Inter. Asso. Theor. Appl. Limnol., 12: 795.
- Wurtz, C. B., 1961. Lethal temperatures for fish: correction of an error. Proc. Pa. Acad. Sci., 35: 195-196.
- Wurtz, C. B., 1962. Zinc effects on fresh water mollusks. Nautilus, 76(2): 53-61.
- Wurtz, C. B., 1962. The effect of heated discharges on aquatic life and water use. ASME Paper No. 61-WA-142, Presented at ASME Mtg., Nov. 26 - Dec. 1, 1961.
- Wurtz, C. B., and C. H. Bridges, 1960. A study of the effects of increased temperatures on biochemical oxygen demand. Proc. Pa. Acad. Sci. 34: 69-73. Bio. Abstr. 36(12), Abstr. No. 36222, 1961. Spo. Fish. Abstr., 6(3), Abstr. No. 4395, 1961.
- Wurtz, C. G., and T. Dolan, 1960. A biological method used in the evaluation of effects of thermal discharge in the Schuylkill River. Purdue Indus. Waste Conf., Engng. Bull. 45(2): 461-472.
- Yamamoto, T., 1931. Temperature constant for the rate of heart beats in Oryzias latipes. Jour. Fac. Sci. Imp. Univ. Tokyo Zool., 2: 381-388.
- Yamamoto, T., 1937. Influence of temperature on the embryonic development of the pond smelt, Hypomesus alidus. Pallas Bull., Jap. Soc. Sci., Fish. Tokyo, 5: 326-332.
- Yamamoto, T., 1937. Influence of temperature on the embryonic development of the European carp Carassius carassius L. Bull. Jap. Soc. Sci. Fish. Tokyo, 5: 326-332.
- Yount, J. L., 1956. Factors that control species numbers in Silver Springs, Florida. Limnol. and Oceano., 1: 286-295.

LABORATORY INVESTIGATIONS SERIES

1. Sargent Polarographic Oxygen Analyzer; Thermo-Fishometer Water Thermometer; Modified Methods for Turbidity; Modified Methods for Color; Suspended Solids Determination (June 24, 1963).*
2. Recovery of Simple Cyanides by the Serfass Distillation Procedure as Compared with the Williams Cuprous Chloride Method (April 7, 1964).
3. Nitrate Determination in Saline and Estuarine Waters (October 13, 1964).
4. Dissolved Oxygen Determinations by Oxygen Meter (January 31, 1966).
5. Picture-Key to the Genera of Aquatic Midges (November, 1967).
6. Temperature and Aquatic Life (December, 1967).

* Out of Print

**LABORATORY INVESTIGATIONS
SERIES — NUMBER SIX**